

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
15 March 2001 (15.03.2001)

PCT

(10) International Publication Number  
**WO 01/18225 A1**

(51) International Patent Classification<sup>1</sup>: C12N 15/90,  
A01K 67/027

(21) International Application Number: PCT/US99/30078

(22) International Filing Date:  
16 December 1999 (16.12.1999)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/152,522 3 September 1999 (03.09.1999) US

(71) Applicant: XENOGEN CORPORATION [US/US]; 860  
Atlantic Avenue, Alameda, CA 94501 (US).

(72) Inventor: ZHANG, Ning; Xenogen Corporation, 860 At-  
lantic Avenue, Alameda, CA 94501 (US).

(74) Agent: SHOLTZ, Charles, K.; Xenogen Corporation,  
860 Atlantic Avenue, Alameda, CA 94501 (US).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,  
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,  
ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,  
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,  
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,  
SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN,  
YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: TARGETING CONSTRUCTS AND TRANSGENIC ANIMALS PRODUCED THEREWITH



(57) Abstract: The present invention teaches targeting constructs and methods of use thereof for creating transgenic animals in which at least one single-copy, non-essential gene is replaced with a reporter expression cassette, for example, a luciferase gene operably linked to a promoter heterologous to the single-copy, non-essential gene. Thus, the present invention provides novel methods and vector constructs useful for the generation of transgenic animals. The invention further includes methods of using these animals.

BEST AVAILABLE COPY

## TARGETING CONSTRUCTS AND TRANSGENIC ANIMALS PRODUCED THEREWITH

5

### TECHNICAL FIELD

This invention is in the field of molecular biology and medicine. More specifically, it relates to novel vector constructs and methods of use thereof for introducing heterologous polynucleotides into a host cell. Further, the invention relates to vector constructs and methods of use thereof to generate transgenic organisms, particularly transgenic mice.

10

### BACKGROUND

In recent years, mouse geneticists have succeeded in creating transgenic animals by manipulating the genes of developing embryos and introducing foreign genes into these embryos. Once these genes have integrated into the genome of the recipient embryo, the resulting embryos or adult animals can be analyzed to determine the function of the gene.

15

Traditionally, transgenic mice have been generated through DNA microinjection approach. Such an approach leads to the creation of "founder" mice having at least one copy of the transgene randomly integrated into the genome. Because neither the copy number nor the integration sites can be controlled, transgenic mice generated by this method are genetically different from each other. The expression of transgenes in such transgenic mice are not uniform because of, for example, the difference in copy numbers of the transgene. Furthermore, chromosomal location of the transgene often affects the expression level of the transgene. For most of *in vivo* studies, particularly ones where it is desirable to compare levels of gene expression across different animals, it is important to have the mice with little or no genetic variation (i.e., isogenic mice) in order to reduce the systematic error.

25

One way of controlling copy number and the chromosomal integration site is by using targeting constructs to create transgenic animals. U.S. Patent Nos. 5,464,764 and 5,487,992 describe this type of transgenic animal in which a gene of interest is deleted or mutated sufficiently to disrupt its function. These "knock-out" animals are made by taking advantage of the phenomena of homologous recombination. (See, also U.S.

30

Patent Nos. 5,631,153 and 5,627,059). Using the "knock-out" approach, when the vector is introduced into the embryonic stem cell, a sequence becomes integrated into a target gene in the genome via homologous recombination. The integration disrupts the function of the target gene allows for examination of the phenotype resulting from the disruption of the gene. Neither of the above methods, however, provide a predictable approach for the generation of gene expression reporter transgenic animals which can be used for quantitative comparisons of gene expression across different animals transfected with the same construct, or for quantitative comparisons of the relative levels of expression of two or more different genes.

10       The present invention solves this and other problems by providing transgenic animals in which at least one single-copy, non-essential gene is replaced with a reporter expression cassette, for example, a luciferase gene operably linked to a heterologous promoter. Thus, the present invention provides novel methods and vector constructs useful for the generation of transgenic animals. The invention further includes methods  
15 of using these animals.

### SUMMARY OF THE INVENTION

The transgenic animals described herein are useful, for example, when studying *in vivo* regulation of selected genes. Also described herein are methods of generating  
20 populations of substantially isogenic transgenic animals, as well as, vectors useful in these methods.

Accordingly, in one embodiment, the subject invention is directed to a transgenic, non-human mammal, for example, a rodent such as a mouse. The mammal comprises at least one single-copy, non-essential gene in its genome, wherein (i) at least a portion of  
25 at least one single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the gene, and (ii) the polynucleotide sequences comprise a first expression cassette which has been introduced into the mammal or an ancestor of the mammal, at an embryonic stage. The first expression cassette typically comprises a first selectable marker, a first transcriptional promoter element heterologous to the gene, and  
30 light-generating protein coding sequences. The light-generating protein coding sequences are operably linked to the promoter element.

The single-copy, non-essential gene may be selected, for example, from the group consisting of vitronectin, *fosB*, and galactin 3 and the first selectable marker may be selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase.

In alternative embodiments, the first transcriptional promoter element is an inducible promoter, a repressible promoter, or a constitutive promoter, and may be selected from the group consisting of VEGF, VEGFR, and TIE2.

In additional embodiments, the transgenic, non-human mammal described above comprises a second single-copy, non-essential gene in its genome, wherein (i) at least a portion of the second single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the second gene, and (ii) the polynucleotide sequences comprise a second expression cassette which has been introduced into the mammal or an ancestor of the mammal, at an embryonic stage. The second expression cassette typically comprises a second selectable marker, a second transcriptional promoter element heterologous to the second gene, and light generating protein coding sequences. The light generating protein coding sequences are operably linked to the promoter element.

The first and second transcriptional promoter elements and selectable markers may be the same or different and the light generating protein in the first expression cassette can produce a different color of light relative to the light generating protein in the second expression cassette.

In yet a further embodiment, the invention is directed to a method of producing a transgenic, non-human mammal, such as a mouse. The mammal has at least one single-copy, non-essential gene in its genome. The method comprises

transfecting an embryonic stem cell of the mammal with a linear vector comprising

(a) a first selectable marker and a reporter expression cassette, the reporter expression cassette comprising a transcriptional promoter element operably linked to a light generating protein coding sequence, and

(b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in said mammal's genome, the targeting polynucleotide sequences



flanking (a), wherein (i) the length of the polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) the transcriptional promoter element is heterologous to the single-copy, non-essential gene;

- 5        selecting embryonic stem cells which each have the first selectable marker and reporter expression cassette integrated into its genome;  
              injecting the embryonic stem cells into a host embryo,  
              implanting the embryo in a foster mother,  
              maintaining the foster mother under conditions which allow production of an  
10    offspring that is a transgenic, non-human mammal carrying the reporter expression cassette.

              In certain embodiments, the offspring is capable of germline transmission of the reporter expression cassette and the method may further comprise breeding the offspring with a mammal which is substantially isogenic with the embryonic stem cells, such that  
15    the breeding yields transgenic F1 offspring carrying the reporter cassette. In particular embodiments, the method comprises breeding the first F1 offspring carrying the reporter cassette with a second F1 offspring carrying the reporter cassette, wherein the breeding yields transgenic F2 offspring carrying the reporter cassette.

              In additional embodiments, the embryonic stems cells may be derived from a  
20    mouse having a dark coat color, the mammal substantially isogenic with the embryonic stem cells may have a light coat color, and/or the F2 offspring carrying the reporter cassette may have a light coat color. In particular embodiments, the embryonic stems cells are derived from a C57BL/6 mouse having a dark coat color, and the mammal substantially isogenic with the embryonic stem cells is a C57BL/6-Tyr C2j/+ mouse  
25    having a light coat color.

              In still a further embodiment, the subject invention is directed to a vector for use in generating a transgenic non-human mammal, for example, a rodent such as a mouse. The mammal has at least one single-copy, non-essential gene in its genome. The vector comprises

- 30        (a) a first selectable marker and a reporter expression cassette, the reporter expression cassette comprising a transcriptional promoter element operably linked to a light generating protein coding sequence, and

(b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in the mammal's genome, the targeting polynucleotide sequences flanking (a), wherein (i) the length of the targeting polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) the transcriptional promoter element is heterologous to the single-copy, non-essential gene.

In certain embodiments, the first selectable marker provides a positive selection and may be selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase. Additionally, the transcriptional promoter element may be an inducible promoter, a repressible promoter, or a constitutive promoter, and may be selected from the group consisting of VEGF, VEGFR, and TIE2.

In alternative embodiments, the vector further comprises a second selectable marker and at least one target polynucleotide sequence is located between the second selectable marker and the first selectable marker. Additionally, the second selectable marker may provide a negative selection and may be selected from the group consisting of adenosine deaminase, thymidine kinase, and dihydrofolate reductase.

The vectors described above may be circular and may contain at least one restriction site whose cleavage results in a linear vector having the following arrangement of elements: target polynucleotide sequence - (a) - targeting polynucleotide sequences or target polynucleotide sequence - (a) - targeting polynucleotide sequences - (second selectable marker).

The coding sequences of the reporter expression cassette present in the vector may comprise codons that are optimal for expression in a host system into which the expression cassette is to be introduced. Additionally, the targeting polynucleotide sequences from single-copy, non-essential genes may be selected from the group consisting of vitronectin, *fosB*, and galactin 3.

The light-generating protein in the mammals and methods described above may be derived from either procaryotic or eucaryotic sources and, in particularly preferred embodiments, the light generating protein is a luciferase.

These and other embodiments of the present invention will be apparent to those of skill in the art in view of the teachings herein.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1 is a schematic depicting construction of the pTK53 vector.

Polynucleotides encoding PGK-P, Neo and TK and 5' and 3' linkers are introduced into  
5 a pKS backbone to produce the vector designated pTK53.

Figure 2 is schematic depicting construction of the pTK-LucR and pTK-LucYG vectors. For pTK-LucR, a polynucleotide encoding LucR is introduced into pTK53. Thus, the pTK-LucR construct contains the PGK-P gene, a neomycin (Neo<sup>r</sup>) gene, a thymidine kinase (TK) gene and sequence encoding red luciferase (Luc-R). For pTK-  
10 LucYG, a polynucleotide encoding LucYG is introduced into pTK53. Thus, the pTK-LucYG construct contains the PGK-P gene, a neomycin (Neo<sup>r</sup>) gene, a thymidine kinase (TK) gene and a sequence encoding yellow-green luciferase (Luc-YG).

Figures 3A is a schematic depicting the vector pTKLR-Vn. Sequences homologous to the vitronectin gene are inserted into pTK-LucR such that they flank the  
15 Neo<sup>r</sup> gene and the Luc-R coding sequence. Figure 3B is a schematic depicting targeting of the linearized pTKLR-Vn vector to the vitronectin chromosomal locus. The VEGF promoter is cloned into the polylinkers between Neo and Luc-R. Upon homologous recombination, the Neo-VEGF-LucR transgene is inserted into the Vn gene. In the figure, (A) shows the targeting vector pTKLR-Vn and (B) shows the mouse vitronectin  
20 gene. In the figure, Neo – neomycin resistance encoding sequences; TK – thymidine kinase encoding sequences; LucR – red luciferase from pGL3Red (Dr. Christopher Contag, Stanford University, Stanford, CA). Regions bearing Vn gene translational start and stop codons are indicated with arrows. Poly(A) sequences are placed upstream of the polylinker to prevent or minimize read-through translation. Figure 3C shows the  
25 nucleotide sequence of vitronectin (SEQ ID NO:38).

Figure 4A is a schematic depicting the vector pTKLG-Fos. Sequences homologous to the FosB gene are inserted into pTK-LucYG such that they flank the Neo<sup>r</sup> gene and the Luc-YG coding sequence. Figure 4B shows the nucleotide sequence of FosB (SEQ ID NO:39).

30 Figure 5A is a schematic depicting targeting of the linearized pTKLG-Fos vector to the FosB chromosomal locus. The VEGFR2 promoter is cloned into the polylinkers between Neo and Luc-YG. Upon homologous recombination, the Neo-VEGFR2-LucYG

transgene will be inserted into a sequence associated with production of FosB. In the figure, (A) shows the targeting vector, and (B) shows the mouse target gene. In the figure, Neo – neomycin resistance encoding sequences; TK – thymidine kinase encoding sequences; LucYG – yellow green luciferase from pGL3-control vector (Promega, Madison, WI). Regions bearing FosB gene translational start and stop codons are indicated with arrows. Poly(A) sequences are placed upstream of the polylinker to prevent or minimize read-through translation. Figure 5B is a schematic depicting targeting of the linearized pTKLG-Fos vector to the FosB chromosomal locus. The TIE2 promoter is cloned into the polylinkers between Neo and Luc-R. Upon homologous recombination, the Neo-Tie2-LucYG transgene is inserted into the FosB gene. In the figure, (A) shows the targeting vector, and (B) shows the mouse target gene. In the figure, Neo – neomycin resistance encoding sequences; TK – thymidine kinase encoding sequences; LucYG – yellow green luciferase from pGL3-control vector (Promega). Regions bearing FosB gene translational start and stop codons are indicated with arrows. Poly(A) sequences are placed upstream of the polylinker to prevent or minimize read-through translation.

Figure 6 depicts PCR conditions for genomic screening for promoters useful in exemplary targeting constructs of the present invention.

Figure 7 depicts generation of targeted transgenic mice using the targeting vectors described herein.

Figure 8 depicts of schematic representation of Southern blot analysis of homologous DNA recombination between pTKLG-Fos targeting vector and the FosB gene.

Figure 9 depicts generation of targeted transgenic mice, using the targeting vectors described herein, and crosses using such transgenics as well as their offspring (F1, first generation; F2, second generation).

Figure 10 depicts crosses using transgenic mice of the present invention to generate dual luciferase transgenic mice.

### MODES FOR CARRYING OUT THE INVENTION

Throughout this application, various publications, patents, and published patent applications are referred to by an identifying citation. The disclosures of these publications, patents, and published patent specifications referenced in this application  
5 are included to more fully describe the state of the art to which this invention pertains.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, cell biology and recombinant DNA, which are within the skill of the art. *See, e.g.*, Sambrook, Fritsch, and Maniatis, MOLECULAR CLONING: A LABORATORY MANUAL, 2nd edition  
10 (1989); CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, (F.M. Ausubel et al. eds., 1987); the series METHODS IN ENZYMOLOGY (Academic Press, Inc.); PCR 2: A PRACTICAL APPROACH (M.J. McPherson, B.D. Hames and G.R. Taylor eds., 1995) and ANIMAL CELL CULTURE (R.I. Freshney. Ed., 1987).

As used in this specification and the appended claims, the singular forms "a,"  
15 "an" and "the" include plural references unless the content clearly dictates otherwise. Thus, for example, reference to "an antigen" includes a mixture of two or more such agents.

#### **Definitions**

20 As used herein, certain terms will have specific meanings.

The terms "nucleic acid molecule" and "polynucleotide" are used interchangeably to and refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Polynucleotides may have any three-dimensional structure, and may perform any function, known or unknown.  
25 Non-limiting examples of polynucleotides include a gene, a gene fragment, exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers.

A polynucleotide is typically composed of a specific sequence of four nucleotide  
30 bases: adenine (A); cytosine (C); guanine (G); and thymine (T) (uracil (U) for thymine (T) when the polynucleotide is RNA). Thus, the term polynucleotide sequence is the alphabetical representation of a polynucleotide molecule. This alphabetical



representation can be input into databases in a computer having a central processing unit and used for bioinformatics applications such as functional genomics and homology searching.

A "coding sequence" or a sequence which "encodes" a selected polypeptide, is a nucleic acid molecule which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide, for example, *in vivo* when placed under the control of appropriate regulatory sequences (or "control elements"). The boundaries of the coding sequence are typically determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence can include, but is not limited to, cDNA from viral, procaryotic or eucaryotic mRNA, genomic DNA sequences from viral or procaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence may be located 3' to the coding sequence. Other "control elements" may also be associated with a coding sequence. A DNA sequence encoding a polypeptide can be optimized for expression in a selected cell by using the codons preferred by the selected cell to represent the DNA copy of the desired polypeptide coding sequence. "Encoded by" refers to a nucleic acid sequence which codes for a polypeptide sequence, wherein the polypeptide sequence or a portion thereof contains an amino acid sequence of at least 3 to 5 amino acids, more preferably at least 8 to 10 amino acids, and even more preferably at least 15 to 20 amino acids from a polypeptide encoded by the nucleic acid sequence. Also encompassed are polypeptide sequences which are immunologically identifiable with a polypeptide encoded by the sequence.

Typical "control elements", include, but are not limited to, transcription promoters, transcription enhancer elements, transcription termination signals, polyadenylation sequences (located 3' to the translation stop codon), sequences for optimization of initiation of translation (located 5' to the coding sequence), translation enhancing sequences, and translation termination sequences. Transcription promoters can include inducible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), repressible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), and constitutive promoters.

"Expression enhancing sequences" typically refer to control elements that improve transcription or translation of a polynucleotide relative to the expression level in the absence of such control elements (for example, promoters, promoter enhancers, enhancer elements, and translational enhancers (e.g., Shine and Delagarno sequences)).

5 "Purified polynucleotide" refers to a polynucleotide of interest or fragment thereof which is essentially free, e.g., contains less than about 50%, preferably less than about 70%, and more preferably less than about 90%, of the protein with which the polynucleotide is naturally associated. Techniques for purifying polynucleotides of interest are well-known in the art and include, for example, disruption of the cell  
10 containing the polynucleotide with a chaotropic agent and separation of the polynucleotide(s) and proteins by ion-exchange chromatography, affinity chromatography and sedimentation according to density.

A "heterologous sequence" as used herein is typically refers to either (i) a nucleic acid sequence that is not normally found in the cell or organism of interest, or (ii) a  
15 nucleic acid sequence introduced at a genomic site wherein the nucleic acid sequence does not normally occur in nature at that site. For example, a DNA sequence encoding a polypeptide can be obtained from yeast and introduced into a bacterial cell. In this case the yeast DNA sequence is "heterologous" to the native DNA of the bacterial cell. Alternatively, a promoter sequence from a Tie2 gene can be introduced into the genomic  
20 location of a *fosB* gene. In this case the Tie2 promoter sequence is "heterologous" to the native *fosB* genomic sequence.

A "polypeptide" is used in it broadest sense to refer to a compound of two or more subunit amino acids, amino acid analogs, or other peptidomimetics. The subunits may be linked by peptide bonds or by other bonds, for example ester, ether, etc. As used  
25 herein, the term "amino acid" refers to either natural and/or unnatural or synthetic amino acids, including glycine and both the D or L optical isomers, and amino acid analogs and peptidomimetics. A peptide of three or more amino acids is commonly called an oligopeptide if the peptide chain is short. If the peptide chain is long, the peptide is typically called a polypeptide or a protein.

30 "Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, a given promoter that is operably linked to a coding sequence (e.g., a reporter expression

cassette) is capable of effecting the expression of the coding sequence when the proper enzymes are present. The promoter or other control elements need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. For example, intervening untranslated yet transcribed sequences can be present between the promoter sequence and the coding sequence and the promoter sequence can still be  
5 considered "operably linked" to the coding sequence.

"Recombinant" as used herein to describe a nucleic acid molecule means a polynucleotide of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation: (1) is not associated with all or a portion of the polynucleotide  
10 with which it is associated in nature; and/or (2) is linked to a polynucleotide other than that to which it is linked in nature. The term "recombinant" as used with respect to a protein or polypeptide means a polypeptide produced by expression of a recombinant polynucleotide. "Recombinant host cells," "host cells," "cells," "cell lines," "cell cultures," and other such terms denoting procaryotic microorganisms or eucaryotic cell  
15 lines cultured as unicellular entities, are used interchangeably, and refer to cells which can be, or have been, used as recipients for recombinant vectors or other transfer DNA, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement to the original parent, due to  
20 accidental or deliberate mutation. Progeny of the parental cell which are sufficiently similar to the parent to be characterized by the relevant property, such as the presence of a nucleotide sequence encoding a desired peptide, are included in the progeny intended by this definition, and are covered by the above terms. An "isolated polynucleotide" molecule is a nucleic acid molecule separate and discrete from the whole organism with  
25 which the molecule is found in nature; or a nucleic acid molecule devoid, in whole or part, of sequences normally associated with it in nature; or a sequence, as it exists in nature, but having heterologous sequences (as defined below) in association therewith.

Techniques for determining nucleic acid and amino acid "sequence identity" also are known in the art. Typically, such techniques include determining the nucleotide  
30 sequence of the mRNA for a gene and/or determining the amino acid sequence encoded thereby, and comparing these sequences to a second nucleotide or amino acid sequence. In general, "identity" refers to an exact nucleotide-to-nucleotide or amino acid-to-amino

acid correspondence of two polynucleotides or polypeptide sequences, respectively.

Two or more sequences (polynucleotide or amino acid) can be compared by determining their "percent identity." The percent identity of two sequences, whether nucleic acid or amino acid sequences, is the number of exact matches between two aligned sequences

5 divided by the length of the shorter sequences and multiplied by 100. An approximate alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman, Advances in Applied Mathematics 2:482-489 (1981). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff, Atlas of Protein Sequences and Structure, M.O. Dayhoff ed., 5 suppl. 3:353-

10 358, National Biomedical Research Foundation, Washington, D.C., USA, and normalized by Gribskov, Nucl. Acids Res. 14(6):6745-6763 (1986). An exemplary implementation of this algorithm to determine percent identity of a sequence is provided by the Genetics Computer Group (Madison, WI) in the "BestFit" utility application. The default parameters for this method are described in the Wisconsin Sequence Analysis

15 Package Program Manual, Version 8 (1995) (available from Genetics Computer Group, Madison, WI). A preferred method of establishing percent identity in the context of the present invention is to use the MPSRCH package of programs copyrighted by the University of Edinburgh, developed by John F. Collins and Shane S. Sturrok, and distributed by IntelliGenetics, Inc. (Mountain View, CA). From this suite of packages

20 the Smith-Waterman algorithm can be employed where default parameters are used for the scoring table (for example, gap open penalty of 12, gap extension penalty of one, and a gap of six). From the data generated the "Match" value reflects "sequence identity." Other suitable programs for calculating the percent identity or similarity between sequences are generally known in the art, for example, another alignment program is

25 BLAST, used with default parameters. For example, BLASTN and BLASTP can be used using the following default parameters: genetic code = standard; filter = none; strand = both; cutoff = 60; expect = 10; Matrix = BLOSUM62; Descriptions = 50 sequences; sort by = HIGH SCORE; Databases = non-redundant, GenBank + EMBL + DDBJ + PDB + GenBank CDS translations + Swiss protein + Spupdate + PIR. Details

30 of these programs can be found at the following internet address:  
<http://www.ncbi.nlm.gov/cgi-bin/BLAST>. When claiming sequences relative to sequences of the present invention, the desired degrees of sequence identity are at least

80%, 85-90%, preferably 92%, more preferably 95%, and even more preferably 98% sequence identity to the reference sequence (i.e., the sequences of the present invention).

Alternatively, the degree of sequence similarity between polynucleotides can be determined by hybridization of polynucleotides under conditions that form stable  
5 duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease(s), and size determination of the digested fragments. Two DNA, or two polypeptide sequences are "substantially homologous" to each other when the sequences exhibit at least about 80%-85%, preferably at least about 85%-90%, more preferably at least about 90%-95%, and most preferably at least about 95%-98%  
10 sequence identity over a defined length of the molecules, as determined using the methods above. As used herein, substantially homologous also refers to sequences showing complete identity to the specified DNA or polypeptide sequence. DNA sequences that are substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that  
15 particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook et al., *supra*; *DNA Cloning, supra*; *Nucleic Acid Hybridization, supra*.

Two nucleic acid fragments are considered to "selectively hybridize" as described herein. The degree of sequence identity between two nucleic acid molecules affects the  
20 efficiency and strength of hybridization events between such molecules. A partially identical nucleic acid sequence will at least partially inhibit a completely identical sequence from hybridizing to a target molecule. Inhibition of hybridization of the completely identical sequence can be assessed using hybridization assays that are well known in the art (e.g., Southern blot, Northern blot, solution hybridization, or the like,  
25 see Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, (1989) Cold Spring Harbor, N.Y.). Such assays can be conducted using varying degrees of selectivity, for example, using conditions varying from low to high stringency. If conditions of low stringency are employed, the absence of non-specific binding can be assessed using a secondary probe that lacks even a partial degree of sequence identity  
30 (for example, a probe having less than about 30% sequence identity with the target molecule), such that, in the absence of non-specific binding events, the secondary probe will not hybridize to the target.



When utilizing a hybridization-based detection system, a nucleic acid probe is chosen that is complementary to a target nucleic acid sequence, and then by selection of appropriate conditions the probe and the target sequence "selectively hybridize," or bind, to each other to form a hybrid molecule. A nucleic acid molecule that is capable of hybridizing selectively to a target sequence under "moderately stringent" typically hybridizes under conditions that allow detection of a target nucleic acid sequence of at least about 10-14 nucleotides in length having at least approximately 70% sequence identity with the sequence of the selected nucleic acid probe. Stringent hybridization conditions typically allow detection of target nucleic acid sequences of at least about 10-14 nucleotides in length having a sequence identity of greater than about 90-95% with the sequence of the selected nucleic acid probe. Hybridization conditions useful for probe/target hybridization where the probe and target have a specific degree of sequence identity, can be determined as is known in the art (see, for example, Nucleic Acid Hybridization: A Practical Approach, editors B.D. Hames and S.J. Higgins, (1985) Oxford; Washington, DC; IRL Press).

With respect to stringency conditions for hybridization, it is well known in the art that numerous equivalent conditions can be employed to establish a particular stringency by varying, for example, the following factors: the length and nature of probe and target sequences, base composition of the various sequences, concentrations of salts and other hybridization solution components, the presence or absence of blocking agents in the hybridization solutions (e.g., formamide, dextran sulfate, and polyethylene glycol), hybridization reaction temperature and time parameters, as well as, varying wash conditions. The selection of a particular set of hybridization conditions is selected following standard methods in the art (see, for example, Sambrook, et al., Molecular Cloning: A Laboratory Manual, Second Edition, (1989) Cold Spring Harbor, N.Y.).

A "vector" is capable of transferring gene sequences to target cells. Typically, "vector construct," "expression vector," and "gene transfer vector," mean any nucleic acid construct capable of directing the expression of a gene of interest and which can transfer gene sequences to target cells. Thus, the term includes cloning, and expression vehicles, as well as integrating vectors.

"Nucleic acid expression vector" or "expression cassette" refers to an assembly which is capable of directing the expression of a sequence or gene of interest. The



nucleic acid expression vector includes a promoter which is operably linked to the sequences or gene(s) of interest. Other control elements may be present as well.

Expression cassettes described herein may be contained within a plasmid construct. In addition to the components of the expression cassette, the plasmid construct may also  
5 include a bacterial origin of replication, one or more selectable markers, a signal which allows the plasmid construct to exist as single-stranded DNA (*e.g.*, a M13 origin of replication), a multiple cloning site, and a "mammalian" origin of replication (*e.g.*, a SV40 or adenovirus origin of replication).

An "expression cassette" comprises any nucleic acid construct capable of  
10 directing the expression of a gene/coding sequence of interest. Such cassettes can be constructed into a "vector," "vector construct," "expression vector," or "gene transfer vector," in order to transfer the expression cassette into target cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

"Luciferase," unless stated otherwise, includes prokaryotic and eukaryotic  
15 luciferases, as well as variants possessing varied or altered optical properties, such as luciferases that produce different colors of light (*e.g.*, Kajiyama, N., and Nakano, E., *Protein Engineering* 4(6):691-693 (1991)).

"Light-generating" is defined as capable of generating light through a chemical reaction or through the absorption of radiation.

20 A "light generating protein" or "light-emitting protein" is a protein capable of generating light in the visible spectrum (between approximately 350 nm and 800 nm). Examples include bioluminescent proteins such as luciferases, *e.g.*, bacterial and firefly luciferases, as well as fluorescent proteins such as green fluorescent protein (GFP).

"Light" is defined herein, unless stated otherwise, as electromagnetic radiation  
25 having a wavelength of between about 300 nm and about 1100 nm.

"Animal" as used herein typically refers to a non-human mammal, including, without limitation, farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs; birds, including domestic, wild and game birds such as chickens,  
30 turkeys and other gallinaceous birds, ducks, geese, and the like. The term does not denote a particular age. Thus, both adult and newborn individuals are intended to be covered.

A "transgenic animal" refers to a genetically engineered animal or offspring of genetically engineered animals. A transgenic animal usually contains material from at least one unrelated organism, such as from a virus, plant, or other animal. The "non-human animals" of the invention include vertebrates such as rodents, non-human  
5 primates, sheep, dogs, cows, amphibians, birds, fish, insects, reptiles, etc. The term "chimeric animal" is used to refer to animals in which the heterologous gene is found, or in which the heterologous gene is expressed in some but not all cells of the animal.

"Analyte" as used herein refers to any compound or substance whose effects (e.g., induction or repression of a specific promoter) can be evaluated using the test  
10 animals and methods of the present invention. Such analytes include, but are not limited to, chemical compounds, pharmaceutical compounds, polypeptides, peptides, polynucleotides, and polynucleotide analogs. Many organizations (e.g., the National Institutes of Health, pharmaceutical and chemical corporations) have large libraries of chemical or biological compounds from natural or synthetic processes, or fermentation  
15 broths or extracts. Such compounds/analytes can be employed in the practice of the present invention.

As used herein, the term "positive selection marker" refers to a gene encoding a product that enables only the cells that carry the gene to survive and/or grow under certain conditions. For example, plant and animal cells that express the introduced  
20 neomycin resistance (Neo<sup>r</sup>) gene are resistant to the compound G418. Cells that do not carry the Neo<sup>r</sup> gene marker are killed by G418. Other positive selection markers will be known to those of skill in the art. Typically, positive selection markers encode products that can be readily assayed. Thus, positive selection markers can be used to determine whether a particular DNA construct has been introduced into a cell, organ or tissue.

25 "Negative selection marker" refers to gene encoding a product which can be used to selectively kill and/or inhibit growth of cells under certain conditions. Non-limiting examples of negative selection inserts include a herpes simplex virus (HSV)-thymidine kinase (TK) gene. Cells containing an active HSV-TK gene are incapable of growing in the presence of gangcylovir or similar agents. Thus, depending on the substrate, some  
30 gene products can act as either positive or negative selection markers.

The term "homologous recombination" refers to the exchange of DNA fragments between two DNA molecules or chromatids at the site of essentially identical nucleotide

sequences. It is understood that substantially homologous sequences can accommodate insertions, deletions, and substitutions in the nucleotide sequence. Thus, linear sequences of nucleotides can be essentially identical even if some of the nucleotide residues do not precisely correspond or align (see, above).

5       A “knock-out” mutation refers to partial or complete loss of expression of at least a portion the target gene. Examples of knock-out mutations include, but are not limited to, gene-replacement by heterologous sequences, gene disruption by heterologous sequences, and deletion of essential elements of the gene (e.g., promoter region, portions of a coding sequence). A “knock-out” mutation is typically identified by the phenotype  
10       generated by the mutation.

      A “single-copy gene” as used herein refers to a gene represented in an organism’s genome only by a single copy at a particular chromosomal locus. Accordingly, a diploid organism has two copies of the gene and both copies occur at the same chromosomal location.

15       A “gene” as used in the context of the present invention is a sequence of nucleotides in a genetic nucleic acid (chromosome, plasmid, etc.) with which a genetic function is associated. A gene is a hereditary unit, for example of an organism, comprising a polynucleotide sequence (e.g., a DNA sequence for mammals) that occupies a specific physical location (a “gene locus” or “genetic locus”) within the  
20       genome of an organism. A gene can encode an expressed product, such as a polypeptide or a polynucleotide (e.g., tRNA). Alternatively, a gene may define a genomic location for a particular event/function, such as the binding of proteins and/or nucleic acids (e.g., phage attachment sites), wherein the gene does not encode an expressed product. Typically, a gene includes coding sequences, such as, polypeptide encoding sequences,  
25       and non-coding sequences, such as, promoter sequences, poly-adenylation sequences, transcriptional regulatory sequences (e.g., enhancer sequences). Many eucaryotic genes have “exons” (coding sequences) interrupted by “introns” (non-coding sequences). In certain cases, a gene may share sequences with another gene(s) (e.g., overlapping genes).

      “Isogenic” means two or more organisms or cells that are considered to be  
30       genetically identical. “Substantially isogenic” means two or more organisms or cells wherein, at the majority of genetic loci (e.g., greater than 99.000%, preferably more than 99.900%, more preferably greater than 99.990%, even more preferably greater than

99.999%), there exists genetic identity between the organisms or cells being compared. In the context of the present invention, two organisms (for example, mice) are considered to be "substantially isogenic" if, for example, inserted transgenes are the primary differences between the genetic make-up of the mice being compared. Further, if, for example, the genetic backgrounds of the mice being compared are the same with the exception that one of the mice has one or several defined mutation(s) (for example, affecting coat color), then these mice are considered to be substantially isogenic. An example of two strains of substantially isogenic mice are C57BL/6 and C57BL/6-Tyr C2j/+.

10       A "pseudogene" as used herein, refers to a type of gene sequence found in the genomes, typically, of eucaryotes, where the sequence closely resembles a known functional gene, but differs in that the pseudogene is non-functional. For example, the pseudogene sequence may contain several stop codons in what would correspond to an open reading frame in the functional gene. Pseudogenes can also have deletions or  
15       insertions relative to their corresponding functional gene. If, for example, in a genome there is a functional gene and a related pseudogene, the functional gene is considered to be a single-copy gene (accordingly, the pseudogene is considered to be single-copy as well).

      A "non-essential gene" refers to a gene whose deletion, disruption, elimination,  
20       reduction of gene function, or mutation is non-lethal, and does not obviously adversely affect the organisms' ability to mature and reproduce. A "non-essential gene with no phenotype" refers to a non-essential gene whose deletion, disruption, elimination, reduction of gene function or mutation has no deleterious effect on the organism. Typically there are no phenotypically reflected gene dosage effects associated with  
25       modification of a non-essential gene with no phenotype -- for example, deletion, disruption or mutation of both copies of a non-essential gene with no phenotype in a diploid organism has essentially the same effect as deletion, disruption, or mutation of one of the two copies present in the diploid organism. In the context of the present invention, a non-essential gene is typically one whose function has been eliminated (e.g.,  
30       by a deletion mutation) and such elimination of function was non-lethal and the organism developed, matured, and was able to reproduce.

The "native sequence" or "wild-type sequence" of a gene is the polynucleotide sequence that comprises the genetic locus corresponding to the gene, e.g., all regulatory and open-reading frame coding sequences required for expression of a completely functional gene product as they are present in the wild-type genome of an organism. The native sequence of a gene can include, for example, transcriptional promoter sequences, translation enhancing sequences, introns, exons, and poly-A processing signal sites. It is noted that in the general population, wild-type genes may include multiple prevalent versions that contain alterations in sequence relative to each other and yet do not cause a discernible pathological effect. These variations are designated "polymorphisms" or "allelic variations."

By "replacement sequence" is meant a polynucleotide sequence that is substituted for at least a portion of the native or wild-type sequence of a gene.

"Linear vector" or "linearized vector," as used herein, is a vector having two ends. For example, circular vectors, such as plasmids, can be linearized by digestion with a restriction endonuclease that cuts at a single site in the plasmid. Preferably, the targeting vectors described herein are linearized such that the ends are not within the targeting sequences.

### General Overview

The present invention relates to vector constructs and methods of creating transgenic animals to be used, for example, as test systems. Methods of using the animals of the present invention include, but are not limited to, using these animals for studies involving tumor growth and other disease conditions. In the practice of the present invention, transgenic, non-human mammals are constructed where a single-copy, non-essential gene is replaced by a reporter expression cassette, preferably a gene encoding a light-generating protein, such as a luciferase-encoding gene, operably linked to a promoter. A variety of promoters are useful in the practice of the present invention, for example, promoters derived from genes associated with tumorigenesis or angiogenesis. Thus, an exemplary promoter can be one that is associated with proteins induced during tumorigenesis, for instance in the presence of tumor generating compounds or of tumors themselves. In this way, expression of the reporter cassette is induced in the animal when, for example, tumors are present, and progression of the

tumor can be evaluated by non-invasive imaging methods using the whole animal. Another exemplary promoter is one that is derived from a gene associated with angiogenesis. Because the promoter is linked to a reporter such as luciferase, non-invasive monitoring of the progression of angiogenesis is possible.

5           Various forms of the different embodiments of the invention, described herein, may be combined.

Non-invasive imaging and/or detecting of light-emitting conjugates in mammalian subjects was described in U.S. Patent No. 5,650,135, by Contag, et al., issued 22 July 1997. This imaging technology can be used in the practice of the present invention in view of the teachings of the present specification. In the imaging method, the conjugates contain a biocompatible entity and a light-generating moiety. Biocompatible entities include, but are not limited to, small molecules such as cyclic organic molecules; macromolecules such as proteins; microorganisms such as viruses, bacteria, yeast and fungi; eukaryotic cells; all types of pathogens and pathogenic substances; and particles such as beads and liposomes. In another aspect, biocompatible entities may be all or some of the cells that constitute the mammalian subject being imaged, for example, cells carrying the vector constructs of the present invention expressing a reporter expression cassette.

Light-emitting capability is conferred on the biocompatible entities by the conjugation of a light-generating moiety. Such moieties include fluorescent molecules, fluorescent proteins, enzymatic reactions giving off photons and luminescent substances, such as bioluminescent proteins. In the context of the present invention, light emitting capability is typically conferred on target cells by having at least one copy of a light-generating protein, e.g., a luciferase, present. In preferred embodiments, luciferase is operably linked to appropriate control elements which can facilitate expression of a polypeptide having luciferase activity. Substrates of luciferase can be endogenous to the cell or applied to the cell or system (e.g., injection into a transgenic mouse, having cells carrying a luciferase construct, of a suitable substrate for the luciferase, for example, luciferin). The conjugation may involve a chemical coupling step, genetic engineering of a fusion protein, or the transformation of a cell, microorganism or animal to express a light-generating protein.



### Targeting Constructs

The targeting cassettes described herein typically include the following components: (1) a suitable vector backbone; (2) a polynucleotide encoding a light generating protein (3) a promoter operably linked to the light generating protein-  
5 encoding gene, wherein the promoter is heterologous to the light generating protein coding sequences; (4) a sequence encoding a positive selection marker; (5) insertion sites flanking the sequence encoding the positive selection marker and the polynucleotide encoding a light generating protein gene, for insertion of sequences which target a single-copy, non-essential chromosomal gene; and, optionally, (6) a sequence encoding a  
10 negative selection marker. Exemplary targeting constructs are shown in Figures 3B, 5A and 5B and described in Examples 1-3.

Suitable vector backbones generally include an F1 origin of replication; a colE1 plasmid-derived origin of replication; polyadenylation sequence(s); sequences encoding antibiotic resistance (*e.g.*, ampicillin resistance) and other regulatory or control elements.  
15 Non-limiting examples of appropriate backbones include: pBluescriptSK (Stratagene, La Jolla, CA); pBluescriptKS (Stratagene, La Jolla, CA) and other commercially available vectors.

In one aspect of the invention the light generating protein is luciferase. Luciferase coding sequences useful in the practice of the present invention include, but  
20 are not limited to, sequences obtained from *lux* genes (procaryotic genes encoding a luciferase activity) and *luc* genes (eucaryotic genes encoding a luciferase activity). A variety of luciferase encoding genes have been identified including, but not limited to, the following: B.A. Sherf and K.V. Wood, U.S. Patent No. 5,670,356, issued 23 September 1997; Kazami, J., et al., U.S. Patent No. 5,604,123, issued 18 February 1997;  
25 S. Zenno, et al, U.S. Patent No. 5,618,722; K.V. Wood, U.S. Patent No. 5,650,289, issued 22 July 1997; K.V. Wood, U.S. Patent No. 5,641,641, issued 24 June 1997; N. Kajiyama and E. Nakano, U.S. Patent No. 5,229,285, issued 20 July 1993; M.J. Cormier and W.W. Lorenz, U.S. Patent No. 5,292,658, issued 8 March 1994; M.J. Cormier and W.W. Lorenz, U.S. Patent No. 5,418,155, issued 23 May 1995; de Wet, J.R., et al,  
30 *Molec. Cell. Biol.* 7:725-737, 1987; Tatsumi, H.N., et al, *Biochim. Biophys. Acta* 1131:161-165, 1992; and Wood, K.V., et al, *Science* 244:700-702, 1989. Eukaryotic luciferase catalyzes a reaction using luciferin as a luminescent substrate to produce light,

whereas prokaryotic luciferase catalyzes a reaction using an aldehyde as a luminescent substrate to produce light.

Wild-type firefly luciferases typically have an emission maxima at about 550 nm. Numerous variants with differing emission maxima have also been studied. For example, Kajiyama and Nakano (*Protein Eng.* 4(6):691-693, 1991; U.S. Patent No. 5,330,906, issued 19 July 1994) teach five variant firefly luciferases generated by single amino acid changes to the *Luciola cruciata* luciferase coding sequence. The variants have emission peaks of 558 nm, 595 nm, 607 nm, 609 nm and 612 nm. A yellow-green luciferase with an emission peak of about 540 nm is commercially available from Promega, Madison, WI under the name pGL3. A red luciferase with an emission peak of about 610 nm is described, for example, in Contag et al. (1998) *Nat. Med.* 4:245-247 and Kajiyama et al. (1991) *Prot. Eng.* 4:691-693.

Positive selection markers include any gene which a product that can be readily assayed. Examples include, but are not limited to, a hprt gene (Littlefield, J. W., Science 145:709-710 (1964)), a xanthine-guanine phosphoribosyltransferase (gpt) gene, or an adenosine phosphoribosyltransferase (aprt) gene (Sambrook et al., *supra*), a thymidine kinase gene (i.e. "TK") and especially the TK gene of herpes simplex virus (Giphart-Gassler, M. et al., *Mutat. Res.* 214:223-232 (1989)), a nptII gene (Thomas, K. R. et al., *Cell* 51:503-512 (1987); Mansour, S. L. et al., *Nature* 336:348-352 (1988)), or other genes which confer resistance to amino acid or nucleoside analogues, or antibiotics, etc, for example, gene sequences which encode enzymes such as dihydrofolate reductase (DHFR) enzyme, adenosine deaminase (ADA), asparagine synthetase (AS), hygromycin B phosphotransferase, or a CAD enzyme (carbamyl phosphate synthetase, aspartate transcarbamylase, and dihydroorotase). Addition of the appropriate substrate of the positive selection marker can be used to determine if the product of the positive selection marker is expressed, for example cells which do not express the positive selection marker nptII, are killed when exposed to the substrate G418 (Gibco BRL Life Technology, Gaithersburg, MD).

The targeting vector typically contains insertion sites for inserting targeting sequences (e.g., sequences that are substantially homologous to the target sequences in the host genome where integration of the targeting vector/expression cassette is desired). These insertion sites are preferably included such that there are two sites, one site on

either side of the sequences encoding the positive selection marker, light generating protein (e.g., luciferase) and the promoter. Insertion sites are, for example, restriction endonuclease recognition sites, and can, for example, represent unique restriction sites. In this way, the vector can be digested with the appropriate enzymes and the targeting  
5 sequences ligated into the vector.

Optionally, the targeting construct can contain a polynucleotide encoding a negative selection marker. Suitable negative selection markers include, but are not limited to, HSV-tk (see, e.g., Majzoub et al. (1996) *New Engl. J. Med.* 334:904-907 and U.S. Patent No. 5,464,764), as well as genes encoding various toxins including the  
10 diphtheria toxin, the tetanus toxin, the cholera toxin and the pertussis toxin. A further negative selection marker gene is the hypoxanthine-guanine phosphoribosyl transferase (HPRT) gene for negative selection in 6-thioguanine.

Exemplary promoters and single-copy, non-essential genes for use in the vector constructs and methods of the present invention are described below.

15

### Promoters

The targeting constructs and transgenic animals described herein contain a sequence encoding a light generating protein, e.g., luciferase, gene operably linked to a promoter. The promoter may be from the same species as the transgenic animal (e.g.,  
20 mouse promoter used in construct to make transgenic mouse) or from a different species (e.g., human promoter used in construct to make transgenic mouse). The promoter can be derived from any gene of interest. In one embodiment of the present invention, the promoter is derived from a gene whose expression is induced during angiogenesis, for example pathogenic angiogenesis like tumor development. Thus, when a tumor begins  
25 to develop in a transgenic animal carrying a vector construct of the present invention, the promoter is induced and the animal expresses luciferase, which can then be monitored *in vivo*.

Exemplary promoters for use in the present invention are selected such that they are functional in a cell type and/or animal into which they are being introduced.

30 Exemplary promoters include, but are not limited to, promoters obtained from the following mouse genes: vascular endothelial growth factor (VEGF) (VEGF promoter described in U.S. Patent No. 5,916,763; Shima et al. (1996) *J. Bio. Chem.* 271:3877-

3883; sequence available on NCBI under accession number U41383); VEGFR2, also known as Flk-1, (VEGFR-2 promoter described, for example, in Röncke et al. (1996) *Circ. Res.* 79:277-285; Patterson et al. (1995) *J. Bio. Chem.* 270:23111-23118; Kappel et al. (1999) *Blood* 93:4282-4292; sequence available as accession number X89777 of NCBI database); Tie2, also known as Tek (Tie2 promoter described, for example, in Fadel et al. (1998) *Biochem. J.* 338:335-343; Schlaeger et al. (1995) *Develop.* 121:1089-1098; Schlager et al. (1997) *PNAS USA* 94:3058-3063). VEGF is a specific mitogen for EC *in vitro* and a potent angiogenic factor *in vivo*. In a tumorigenesis study, it was shown that VEGF was critical for the initial subcutaneous growth of T-47D breast carcinoma cells transplanted into nude mice, whereas other angiogenic factors, such as, bFGF can compensate for the loss of VEGF after the tumors have reached a certain size (Yoshiji, H., et al., 1997 *Cancer Research* 57: 3924-28). VEGF is a major mediator of aberrant EC proliferation and vascular permeability in a variety of human pathologic situation, such as, tumor angiogenesis, diabetic retinopathy and rheumatoid arthritis (Benjamin LE, et al., 1997 *PNAS* 94: 8761-66; Soker, S., et al., 1998 *Cell* 92: 735-745). VEGF is synthesized by tumor cells *in vivo* and accumulates in nearby blood vessels. Because leaky tumor vessels initiate a cascade of events, which include plasma extravasation and which lead ultimately to angiogenesis and tumor stroma formation, VEGF plays a pivotal role in promoting tumor growth (Dvorak, H.F., et al., 1991 *J Exp Med* 174:1275-8). VEGF expression was upregulated by hypoxia (Shweiki, D., et al., 1992 *Nature* 359: 843-5). VEGF is also upregulated by overexpression of v-Src oncogene (Mukhopadhyay, D., et al., 1995 *Cancer Res.* 15: 6161-5), c-SRC (Mukhopadhyay, D., et al., 1995 *Nature* 375: 577-81), and mutant ras oncogene (Plate, K.H., et al., 1992 *Nature* 359: 845-8). The tumor suppressor p53 downregulates VEGF expression (Mukhopadhyay, D., et al., 1995 *Cancer Res.* 15: 6161-5).

A number of cytokines and growth factors, including PGF and TPA (Grugel, S., et al., 1995 *J. Biological Chem.* 270: 25915-9), EGF, TGF- $\beta$ , IL-1, IL-6 induce VEGF mRNA expression in certain type of cells (Ferrara, N., et al., 1997 *Endocr. Rev.* 18: 4-25). Kaposi's sarcoma-associated herpesvirus (KSHV) encoded a G-protein-coupled receptor, a homolog of IL-8 receptor, can activate JNK/SAPK and p38MAPK and increase VEGF production, thus causing cell transformation and tumorigenicity (Bais, C., et al., *Nature* 1998 391:86-9). VEGF overexpression in skin of transgenic mice

induces angiogenesis, vascular hyperpermeability and accelerated tumor development (Larcher, F., et al., *Oncogene* 1998 17:303-11).

VEGF-B (cDNA sequences available on databases) is a mitogen for EC and may be involved in angiogenesis in muscle and heart (Olofsson, B., et al., 1996 *Proc Natl Acad Sci U S A* 93:2576-81). Shown *in vitro*, binding of VEGF-B to its receptor VEGFR-1 leads to increased expression and activity of urokinase type plasminogen activator and plasminogen activator inhibitor, suggesting a role for VEGF-B in the regulation of extracellular matrix degradation, cell adhesion, and migration (Olofsson, B., et al., 1998 *Proc Natl Acad Sci U S A* 95:11709-14).

10 VEGF-C (see, e.g., U.S. Patent No. 5,916,763 and Shima et al., *supra*) may regulate angiogenesis of lymphatic vasculature, as suggested by the pattern of VEGF-C expression in mouse embryos (Kukk, E., et al., 1996 *Development* 122: 3829-37). Although VEGF-C is also a ligand for VEGFR-2, the functional significance of this potential interaction is unknown. Overexpression of VEGF-C in the skin of transgenic mice resulted in lymphatic, but not vascular, endothelial proliferation and vessel enlargement, suggesting the major function of VEGF-C is through VEGFR-3 rather than VEGFR-2 (Jeltsch M, et al., 1997 *Science* 276:1423-5). Shown by the CAM assay, VEGF and VEGF-C are specific angiogenic and lymphangiogenic growth factors, respectively (Oh, S.J., et al., (1997) *Devel. Biol.* 188: 96-109). VEGF-C overexpression 15 in the skin of transgenic mice resulted in lymphatic, but not vascular, endothelial proliferation and vessel enlargement (Jeltsch M, et al., 1997 *Science* 276:1423-5).

VEGF-D (cDNA sequences available on databases) is a mitogen for EC. Given that VEGF-D can also activate VEGFR-3, it is possible that VEGF-D could be involved in the regulation of growth and/or differentiation of lymphatic endothelium (Achen, M.G., et al., 1998 *Proc Natl Acad Sci U S A* 95: 548-53). VEGF-D is induced by transcription factor c-Fos in mouse (Orlandini, M., 1996 *PNAS* 93: 11675-80).

VEGFR-1 signaling pathway may regulate normal endothelial cell-cell or cell matrix interactions during vascular development, as suggested by the knockout study (Fong, G.H., et al., 1995 *Nature* 376: 65-69). Although VEGFR-1 has a higher affinity 30 to VEGF than VEGFR-2, it does not transduce the mitogenic signals of VEGF in ECs (Soker, S., et al., 1998 *Cell* 92: 735-745). VEGFR-2 (see, e.g., Röncke et al., Patterson et al., Kappel et al. (1999), *supra*) appears to be the major transducer of VEGF signals in



EC that result in chemotaxis, mitogenicity and gross morphological changes in target cells (Soker, S., et al., 1998 *Cell* 92: 735-745). VEGFR-3 has an essential role in the development of the embryonic cardiovascular system before the emergence of lymphatic vessels, as shown by the knockout study (Dumont, D.J., et al., 1998 *Science* 282: 946-949). Neuropilin-1 (see, e.g., Soker et al. (1998) *Cell* 92:735-745) is a receptor for VEGF165. It can enhance the binding of VEGF165 to VEGFR-2 and VEGF165 mediated chemotaxis (Soker, S., et al., 1998 *Cell* 92: 735-745). Neuropilin1 overexpression in transgenic mice resulted in embryonic lethality. The embryos possessed excess capillaries and blood vessels. Dilated vessels and hemorrhage were also observed (Kitsukawa, T., et al., 1995 *Development* 121: 4309-18).

Further promoters of interest include, but are not limited to, the following. Ang2 is expressed only at predominant vascular remodeling sites, such as ovary, placenta, uterus (Maisonpierre, P.C., et al., 1997 *Science* 277: 55-60). In glioblastoma angiogenesis, Ang2 is found to be expressed in endothelial cells of small blood vessel and capillaries while Ang1 is expressed in glioblastoma tumor cells (Stratmann, A., 1998 *Am J Pathol* 153: 1459-66). Ang2 is up-regulated in bovine microvascular endothelial by VEGF, bFGF, cytokines, hypoxia (Mandriota, S.J., 1998 *Circ Res* 83: 852-9). Ang2 transgenic overexpression disrupts angiogenesis, and is embryonic lethal (Maisonpierre, P.C., et al., 1997 *Science* 277: 55-60). Ang1 is widely expressed, less abundant in heart and liver (Maisonpierre, P.C., et al., 1997 *Science* 277: 55-60). Ang1 is expressed in mesenchymal cells and may up-regulate the expression of Tie2 in the endothelial cells (Suri, C., et al., 1996 *Cell* 87: 1171-1180). Ang1 overexpression in the skin of transgenic mice produces larger, more numerous, and more highly branched vessels (Suri, C., et al., *Science* 1998 282:468-71). Tie2 (see, e.g., Fadel et al.; Schlaeger et al. (1995), and Schlaeger et al. (1997), *supra*) is endothelial cell specific, up-regulated during wound healing, follicle maturation (Puri, M.C., et al., 1995 *EMBO J* 14: 5884-91) and pathologic angiogenesis (Kaipainen, A., 1994 *Cancer Research* 54: 6571-77), such as, glioblastoma (Stratmann, A., 1998 *Am J Pathol* 153: 1459-66). Tie2 is also expressed in non-proliferating adult endothelium and endothelial cell lines (Dumont, D.J., et al. (1994) *Genes & Develop.* 8:1897-1909). A Tie2 activating mutation causes vascular dysmorphogenesis (Vikkula M, et al., 1996 *Cell* 87: 1181-1190). Tie2 mutant overexpression in transgenic mice is embryonic lethal (Dumont, D.J., et al., *supra*).



Other promoters useful in the practice of the present invention include, by way of example, promoters derived from the sequences encoding the following polypeptide products: PTEN (dual specificity phosphatase); BAI (brain-specific angiogenesis inhibitor); KAI1 (KANGAI 1); catenin beta-1 (cadherin-associated protein, beta); COX2 (PTGS2 cyclooxygenase 2, a.k.a. prostaglandin-endoperoxide synthase 2); MMP2 (72 kDa Type IV-A collagenase); MMP9 (92 kDa type IV-B collagenase); TIMP2 (tissue inhibitor of metalloproteinase 2); and TIMP3 (tissue inhibitor of metalloproteinase 3).

PTEN is a tumor suppressor gene and encodes a protein of 403 amino acids. (Li et al. (1997) *Science* 275:1943-1946; DiCristofano et al. (1998) *Nature Genet.* 19:348-355). Overexpression of PTEN has been shown to inhibit cell migration and it is postulated that this protein may function as a tumor suppressor by negatively regulating cell interactions with the extracellular matrix or by negatively regulating the PI3K/PKB/Akt signaling pathway. (Tamura et al. (1998) *Science* 280:1614-1617; Stambolic et al. (1998) *Cell* 95:29-29). Mutations in PTEN have been detected in cancer cell lines and in the germline of patients having Cowden disease, Lhermitte-Duclos disease and Bannayan-Zonana syndrome (diseases and syndromes which are characterized by hyperplastic/dysplastic changes in the prostate, skin and colon and which are associated with an increased risk of certain cancers, for example, breast cancer, prostate cancer and colon cancer). (Marsh et al. (1998) *Hum. Molec. Genet.* 7:507-515; Marsh et al. (1998) *J. Med. Genet.* 35:881-885; Nelen et al. (1997) *Hum. Molec. Genet.* 6:1383-1387).

BAI1 protein is predicted to be 1,584 amino acids in length and includes an extracellular domain, an intracellular domain and a 7-span transmembrane region similar to that of the secretin receptor. (Nishimori et al. (1997) *Oncogene* 15:245-2150). The extracellular region of BAI1 has a single Arg-Gly-Asp (RGD) motif recognized by integrins and also has five sequences corresponding to the thrombospondin type I (accession number 188060) repeats that can inhibit angiogenesis includes by basic fibroblast growth factor (bFGF, accession number 134920). Shiratsuchi et al. (1997) *Cytogenet. Cell Genet.* 79:103-108, cloned 2 other brain-specific angiogenesis inhibiting genes, designated BAI2 (accession number 602683) and BAI 3 (accession number 602684). Thus, it is postulated that members of this gene family may play a role in suppression of glioblastoma.

KAI1 encodes a 267 amino acid protein which is a member of the leukocyte surface glycoprotein family. The protein has 4 hydrophobic transmembrane domains and 1 large extracellular hydrophilic domain with three potential N-glycosylation sites. (Dong et al. (1995) *Science* 268:884-886). Molecular analysis of KAI1 is described, for example, in Dong et al. (1997) *Genomics* 41:25-32. KAI1 is a tumor metastasis suppressor gene that is capable of inhibiting the metastatic process in experimental animals. Expression of KAI1 is downregulated during tumor progression of prostate, breast, lung, bladder and pancreatic cancers in humans, apparently at the transcriptional or postranscriptional level. Mashimo et al. (1998) *PNAS USA* 95:11307-11311, found that the tumor suppressor gene p53 can directly inactivate the KAI1 gene by interacting with the region 5' to the coding sequence, suggesting a direct relationship between p53 and KAI1.

Catenin beta-1 is an adherens junction (AJ) protein, which are critical for establishing and maintaining epithelial cell layers, for instance during embryogenesis, wound healing and tumor cell metastasis. Molecular analysis, including description of sequence homology to plakoglobin (accession number 173325), homology to the drosophila gene "armadillo" and interactions with Lef1/Tcf DNA binding proteins, is described, for example, in Nollet et al. (1996) *Genomics* 32:413-424; McCrea et al. (1991) *Science* 254:1359-1361 and Korinek et al. (1997) *Science* 275:1784-1787. In addition, studies by Korinek et al., *supra* and Morin et al. (1997) *Science* 275:1787-1790, have indicated that APC (accession number 175100) negatively regulates catenin beta and that regulation of this protein is critical to the tumor suppressive effect of APC. Abnormally high levels of beta-catenin have been detected in certain human melanoma cell lines. (Rubinfeld et al. (1997) *Science* 275:1790-1792. Koch et al. (1999) *Cancer Res.* 59:269-273 report that childhood hepatoblastomas frequently carry a mutated degradation targeting box of the beta-catenin gene. Transgenic mice which express catenin beta under the control of an epidermal promoter undergo de novo hair morphogenesis and eventually these animals develop two types of tumors – epithelioid cysts and trichofolliculomas. Gat et al. (1998) *Cell* 95:605-614.

COX2 encodes a cyclooxygenase and is a key regulator of prostaglandin synthesis. (Hla et al. (1992) *PNAS USA* 89:7384-7388; Jones et al. (1993) *J. Biol. Chem.* 268:9049-9054). In particular, COX2 is generally considered to be a mediator of

inflammation and overexpression of COX2 in rat epithelial cells results in elevated levels of E-cadherin and Bcl2. (Tsuji & DuBois (1995) *Cell* 83:493-501). In co-cultures of endothelial cells and colon carcinoma cells, cells that overexpress COX2 produce prostaglandins, proangiogenic factors and stimulate both endothelial migration and tube formation. (Tsuji et al. (1998) *Cell* 93:705-716). Experiments conducted using APC knock-out mice have demonstrated that animals homozygous for a disrupted COX2 locus develop significantly more adenomatous polyps. (Oshima et al. (1996) *Cell* 87:803-809). COX-2 "knock out" mice develop severe nephropathy, are susceptible to peritonitis, exhibit reduced arachidonic acid-induced inflammation and exhibit reduced indomethacin-induced gastric ulceration. (Morham et al. (1995) *Cell* 83:473-482; Langenbach et al. (1995) *Cell* 83:483-492). Female mice that are deficient in cyclooxygenase 2 exhibit multiple reproductive failures. (Lim et al. (1997) *Cell* 91:197-208).

MMP2 is a metalloproteinase that specifically cleaves type IV collagen. A C-terminal fragment of MMP2, termed PEX, prevents normal binding to alpha-V/beta-3 and disrupts angiogenesis and tumor growth. (Brooks et al. (1998) *Cell* 92:391-400).

MMP9 is a collagenase secreted from normal skin fibroblasts. MMP9 null mice exhibit an abnormal pattern of skeletal growth plate vascularization and ossification. (Vu et al. (1998) *Cell* 93:411-422).

TIMP2 is a collagenase and appears to play a major role in modulating the activity of interstitial collagenase and a number of connective tissue metalloendoproteases. (Stetler-Stevenson et al. (1989) *J. Biol. Chem.* 264:17372-17378). Unlike TIMP1 and TIMP3, TIMP2 is not upregulated by TPA or TGF-beta. (Hammani et al. (1996) *J. Biol. Chem.* 271:25498-25505).

TIMP3 (Wilde et al. (1994) *DNA Cell Biol.* 13:711-718) is localized in the extracellular matrix in both its glycosylated and unglycosylated forms. Studies of mutant TIMP3 proteins have demonstrated that C-terminal truncations do not bind to the extracellular matrix. (Langton et al. (1998) *J. Biol. Chem.* 273:16778-16781).

As one of skill in the art will appreciate in view of the teachings of the present specification, promoter sequences can be easily derived and isolated from known polypeptide sequences or from cDNA or genomic sequences, using method known in the art in view of the teachings herein. An exemplary method of isolating promoter

sequences using cDNA is via a GenomeWalker® kit, commercially available from Clontech (Palo Alto, CA), and described on page 27 of the 1997-1998 Clontech catalog.

### **Targeting Sequences: Non-Essential Genes**

5        Central to the present invention is the fact that the targeting constructs contain "targeting" sequences (flanking, for example, the light generating protein-encoding sequence and promoter) derived from a single-copy, non-essential gene. These targeting sequences in the construct act via homologous recombination to replace at least a portion of the non-essential gene in the genome with the light-generating protein-encoding (*e.g.*,  
10    luciferase-encoding) sequence operably linked to a promoter.

Non-limiting examples of targeting sequences for use in generating transgenic mice include sequences obtained from or derived from vitronectin, Fos B and galactin 3. A search of Mouse Knockout & Mutation Database (Genome Systems, Inc., St. Louis, MO) can be used to identify genes that have been knocked-out in mice where the  
15    generated knockout mice displayed no obvious defects. The chromosomal locus for all these genes can be used to target promoter-(light generating protein, *e.g.*, luciferase) transgenes similar to what is described in Example 2. Single-copy, non-essential mouse genes identified in this manner include, but are not limited to, the following: Moesin (Msn), Doi Y., et al., J Biol Chem 1999, 274:2315–2321; Plasminogen activator  
20    inhibitor, type II (Planh2) and Planh1, Dougherty K.M., Proc Natl Acad Sci USA 1999, 96:686–691; Protein tyrosine phosphatase, receptor type, B (Ptpnb), Elchebly et al. (1999) Science 283:1544–1548; Presenilin 1 (Psen1), Guo Q, et al. (1999) Proc Natl Acad Sci USA, 96:4125–4130; Protein kinase, mitogen-activated 9 (Prkm9) / SAPK/Erk/kinase 2 (Serk2), Kuan CY et al. (1999) Neuron 4:667–676; CD152 antigen  
25    (Cd152) / CD86 antigen (Cd86) / CD80 antigen (Cd80), Mandelbrot DA, et al. (1999) J Exp Med, 189:435–440; Poly (ADP-ribose) polymerase (Adprp), Masutani M, et al. (1999), Proc Natl Acad Sci USA 96:2301–2304; Sodium channel, nonvoltage-gated 1 beta (Scnn1b), Pradervand S, et al. (1999) Proc Natl Acad Sci USA 96:1732–1737; Nuclear receptor coactivator 1 (Ncoa1), Qi C, et al. (1999) Proc Natl Acad Sci USA  
30    96:1585–1590; Decay accelerating factor 1 (Daf1), Sun X, et al. (1999) Proc Natl Acad Sci USA 1999, 96:628–633; Necdin (Ndn), Tsai TF, et al. (1999) Nat Genet 22:15–16; Relaxin (Rln); Zhao L, et al. (1999) Endocrinology 140:445–453; Adenylyl cyclase 8

- (Adcy8), Abdel-Majid RM, et al. (1998) *Nat Genet* 19:289-291; Leukemia inhibitory factor (Lif), Bugga L, et al. (1998) *J Neurobiol* 36:509-524; Lectin, galactose binding, soluble 3 (Lgals3) and Lgals1, Calnot C, et al. (1998) *Dev Dyn* 211:306-313; Urokinase plasminogen activator receptor (Plaur) Carmeliet P, et al. (1998) *J Cell Biol* 140:233-245; Nitric oxide synthase 1, neuronal (Nos1), Chao DS, et al. (1998) *J Neurochem* 71:784-789; Homeo box A7 (Hoxa7), Chen F, et al. (1998) *Mech Dev* 77:49-57; Myosin light chain, phosphorylatable, cardiac ventricles (Myhpc) Chen J, et al. (1998) *J Biol Chem* 273:1252-1256; Homeo box B7 (Hoxb7), Chen F, et al. (1998) *Mech Dev* 77:49-57; Nuclear factor of kappa light chain gene enhancer in B-cells inhibitor, alpha and beta (Nfkb1a and Nfkb1b), Cheng JD, et al. (1998) *J Exp Med* 186:1055-1062; Enolase 1, alpha non-neuron (Eno1), Couldrey C, et al. (1998) *Dev Dyn* 212:284-292; Xeroderma pigmentosum, complementation group A (Xpa), De Vries A, et al. (1998) *Exp Eye Res* 1998, 67:53-59; Von Willebrand factor homolog (Vwf), Denis C, et al. (1998) *Proc Natl Acad Sci USA* 95:9524-9529; Lysosomal acid lipase 1 (Lip1), Du H, et al. (1998) *Hum Mol Genet* 7:1347-1354; UNC-5 homolog (*C. elegans*) 3 (Unc5h3), Eisenman LM, et al. (1998) *J Comp Neurol* 394:106-117; Protein phosphatase 1, regulatory (inhibitor) subunit 1B (Ppp1r1b), Fienberg AA, et al. (1998) *Science* 281:838-842; Myelin-associated glycoprotein (Mag) Fujita N, et al. (1998) *J Neurosci* 18:1970-1978; Paraoxonase 1 (Pon1), Furlong CE, et al. (1998) *Neurotoxicology* 19:645-650; Brain derived neurotrophic factor (Bdnf), Garek RR, et al. (1998) *Laryngoscope* 108:671-678; Neurotrophin 3 (Ntf3), Garek RR, et al. (1998) *Laryngoscope* 108:671-678; Myoglobin (Mb), Garry DJ, et al. (1998) *Nature* 395:905-908; Opioid receptor, mu (Oprm), Gaveriaux-Ruff C, et al. (1998) *Proc Natl Acad Sci USA* 95:6326-6330; Neuropeptide Y (Npy), Hollopeter G, et al. (1998) *Int J Obes Relat Metab Disord* 22:506-512; Procollagen, type I, alpha 1 (Col1a1) Hormuzdi SG, et al. (1998) *Mol Cell Biol* 18:3368-3375; Centromere autoantigen B (Cenpb), Hudson DF, et al. (1998) *J Cell Biol* 141:309-319; Oculocerebrorenal syndrome of Lowe (ocrl), Janne PA, et al. (1998) *J Clin Invest* 101:2042-2053; arachidonate 12-lipoxygenase (Alox12) Johnson EN, et al. (1998) *Proc Natl Acad Sci USA* 95:3100-3105; H19 fetal liver mRNA (H19), Jones BK, et al. (1998) *Genes Dev* 12:2200-2207; Hepatocyte nuclear factor 3 gamma (winged helix transcription factor) (Hnf3g), Kaestner KH, et al. (1998) *Mol Cell Biol* 18:4245-4251; Bone morphogenetic protein 2 (Bmp2) / Bone



- morphogenetic protein 7 (Bmp7) and Bmp5, Katagiri T, et al. (1998) *Dev Genet* 22:340–348; Intercellular adhesion molecule (Icam1), Ley K, et al. (1998) *Circ Res* 83:287–294; Glutamyl aminopeptidase (Enpep), Lin Q, et al. *J Immunol* 1998, 160:4681–4687; Prion protein (Prnp), Lipp HP, et al. *Behav Brain Res* 1998, 95:47–54; RAB3A, member RAS oncogene family (Rab3a), Lonart G, et al. *Neuron* 1998, 21:1141–1150; Potassium voltage gated channel, shaker related subfamily, member 4 (Kcna4), London B, et al., *J Physiol (Lond)* 1998, 509:171–182; Apurinic/aprimidinic endonuclease (Apex), Ludwig DL, et al. *Mutat Res* 1998, 409:17–29; T-cell receptor gamma, variable 5 (Tcrg-V5), Mallick-Wood CA, et al. *Science* 1998, 279:1729–1733; Nuclear, factor, erythroid derived 2, like 2 (Nfe2l2), Martin F, et al. *Blood* 1998, 91:3459–3466; Interleukin 13 (Il13), McKenzie GJ, et al. *Curr Biol* 1998, 8:339–340; Sorbitol dehydrogenase 1 (Sdh1), Ng T, et al. *Diabetes* 1998, 47:961–966; Guanine nucleotide binding protein, alpha 11 (Gnal1), Offermanns S, et al. *EMBO J* 1998, 17:4304–4312; Estrogen receptor alpha (Estra), Ogawa S, et al. *Endocrinology* 1998, 139:5070–5081; Integrin beta 2 (Itgb2), Intracellular adhesion molecule (Icam1) and CD34 antigen, Oliveira-dos-Santos AJ, et al. *Eur J Immunol* 1998, 28:2882–2892; Angiotensin receptor 1b (Agtr1b), Oliverio MI, et al. *Proc Natl Acad Sci USA* 1998, 95:15496–15501; Complement factor B (factor B), Pekna M, et al. *Scand J Immunol* 1998, 47:375–380; Centromere autoantigen B (Cenpb), Perez-Castro AV, et al. *Dev Biol* 1998, 201:135–143; Procollagen, type V, alpha 2 (Col5a2) / Fibrillin 1 (Fbn1), Phelps RG, et al. *Mol Med* 1998, 4:356–360; Plasminogen activator inhibitor, type 1 (Plnh1), Pinsky DJ, et al. *J Clin Invest* 1998, 102:919–928, Carmeliet P, et al. *J Clin Invest* 1993, 92:2756–2760; Placentae and embryos oncofetal gene (Pem), Pitman JL, et al. *Dev Biol* 1998, 202:196–214; Postmeiotic segregation increased 1 (*S. cerevisiae*) (Pms1), Prolla TA, et al. *Nat Genet* 1998, 18:276–278; Prion protein, structural locus (Prn-p), Prusiner SB, et al. *Proc Natl Acad Sci USA* 1998, 95:10608–10612, Lledo P-M, et al. *Proc Natl Acad Sci U S A* 1996, 93:2403–2407, Sailer A, et al. *Cell* 1994, 77:967–968, Weissmann C, et al. *Philos Trans R Soc Lond [Biol]* 1994, 343:431–433, Bueler H, et al. *Cell* 1993, 73:1337–1347, Weissmann C, et al. *Intervirology* 1993, 35:164–175; NAD (P)H:quinone oxidoreductase, Radjendirane V, et al. *J Biol Chem* 1998, 273:7382–7389; Alpha tropomyosin (Tpm1), Rethinasamy P, et al. *Circ Res* 1998, 82:116–123; Goosecoid and Goosecoid-like (Gsc1), Wakamiya M, et al. *Hum Mol Genet* 1998, 7:1835–1840, Saint-Pore B, et al. *Hum Mol Genet* 1998,



- 7:1841–1849; Schlafen 1 (Slfn1); Schwarz DA, et al. *Immunity* 1998, 9:657–668; Nuclear factor, erythroid derived 2, ubiquitous (Mafk), Shavit JA, et al. *Genes Dev* 1998, 12:2164–2174; Microphthalmia-associated transcription factor (Mitf), Smith SB, et al. *Exp Eye Res* 1998, 66:403–410; Bone morphogenetic protein 6 (Bmp6), Solloway MJ, et al. *Dev Genet* 1998, 22:321–339; Phosphatidylinositol glycan, class A (Piga), Takahama Y, et al. *Eur J Immunol* 1998, 28:2159–2166; Paired-related homeobox 2 (Prx2), Berge D, et al. *Development* 1998, 125:3831–3842; Prostaglandin E receptor EP1 subtype (Ptgerp1), Ushikubi F, et al. *Nature* 1998, 395:281–284; Immunoglobulin kappa chain complex (Igk), van der Stoep N, et al. *Immunity* 1998, 8:743–750; Adenine
- 10 phosphoribosyl transferase (Aprt), Van Sloun PP, et al. *Nucleic Acids Res* 1998, 26:4888–4894; Microtubule associated protein 4 (Mtap4), Voss AK, et al. *Dev Dyn* 1998, 212:258–266; ; 3-hydroxy-3-methylglutaryl-coenzyme A lyase (Hmgcl), Wang SP, et al. *Hum Mol Genet* 1998, 7:2057–2062; Fibroblast growth factor receptor 4 (Fgfr4), Weinstein M, et al. *Development* 1998, 125:3615–3623; Hepsin (Hpn), Wu Q,
- 15 et al. *J Clin Invest* 1998, 101:321–326; Small inducible cytokine A11 (Scya11), Yang T, et al. *Blood* 1998, 92:3912–3923; Small nuclear ribonucleoprotein N (Snrpn), Yang T, et al. *Nat Genet* 1998, 19:25–31; DNA fragmentation factor, alpha subunit (Dffa), Zhang J, et al. *Proc Natl Acad Sci USA* 1998, 95:12480–12485; Early growth response 1 (Egr1), Zheng D, et al. *Neuroscience* 1998, 83:251–258; Early growth response 1 (Egr1) /
- 20 Hormone receptor (Hmr), Zheng D, et al. *Neuroscience* 1998, 83:251–258; Hemochromatosis (Hfe), Zhou XY, et al. *Proc Natl Acad Sci USA* 1998, 95:2492–2497; Alpha tropomyosin (Tpm1), Blanchard EM, et al. *Circ Res* 1997, 81:1005–1010; tRNA phosphoserine (Trsp), Bosl MR, et al. *Proc Natl Acad Sci U S A* 1997, 94:5531–5534; Angiotensin receptor 1b (Agtr1b), Chen X, et al. *Am J Physiol* 1997, 272:F299–F304;
- 25 Xeroderma pigmentosum, complementation group C (Xpc), Cheo DL, et al. *Mut Res* 1997, 374:1–9; B cell leukemia/lymphoma 6 (Bcl6), Dent AL, et al. *Science* 1997, 276:589–592; Fumarylacetoacetate hydrolase (Fah) / 4-hydroxyphenylpyruvic acid dioxygenase (Hpd), Endo F, et al. *J Biol Chem* 1997, 272:24426–24432; N-methylpurine-DNA glycosylase (Mpg), Engelward BP, et al. *Proc Natl Acad Sci USA*
- 30 1997, 94:13087–13092; Interleukin 1 receptor, type 1 (Il1r1), Glaccum MB, et al. *J Immunol* 1997, 159:3364–3371; N-methylpurine-DNA glycosylase (Mpg), Hang B, et al. *Proc Natl Acad Sci USA* 1997, 94:12869–12874; Gamma-aminobutyric acid (GABA-

- A) receptor, subunit alpha 6 (Gabra6), Homanics GE, et al. Mol Pharmacol 1997, 51:588–596; Superoxide dismutase 2, mitochondrial (Sod2), Huang TT, et al. Arch Biochem Biophys 1997, 344:424–432; Interleukin 11 receptor, alpha chain 1 (Il11ra1), Nandurjar HH, et al. Blood 1997, 90:2148–2159; Alkaline phosphatase 5 (Akp5),
- 5 Narisawa S, et al. Dev Dyn 1997, 208:432–446; GATA-binding protein 4 (Gata4), Narita N, et al. Development 1997, 124:3755–3764; Lymphocyte protein tyrosine kinase (Lck) / Fyn protooncogene (Fyn); Page ST, et al. Eur J Immunol 1997, 27:554–562; P glycoprotein 3 (Pgy3), Schinkel AH, et al. Proc Natl Acad Sci U S A 1997, 94:4028–4033; P glycoprotein 1 (Pgy1) / P glycoprotein 3 (Pgy3), Schinkel AH, et al., Proc Natl
- 10 Acad Sci U S A 1997, 94:4028–4033; Creatine kinase, mitochondrial 1, ubiquitous (Ckmt1), Steeghs K, et al. J Neurosci Methods 1997, 71:29–41; T cell receptor gamma, variable 4 (Tcrg-V4), Sunaga S, et al. J Immunol 1997, 158:4223–4228; Ia-associated invariant chain (p31 form) (Ii), Takaesu NT, et al. J Immunol 1997, 158:187–199; Solute carrier family 18 (vesicular monoamine), member 2 (Slc18a2), Takahashi N, et al. Proc
- 15 Natl Acad Sci USA 1997, 94:9938–9943; Matrix metalloproteinase 7 (Mmp7), Wilson CL, et al. Proc Natl Acad Sci U S A 1997, 94:1402–1407; Formin (Fmn), Wynshaw-Boris A, et al. Mol Med 1997, 3:372–384; Synaptophysin (Syp), Arrandale JM, et al. J Biol Chem 1996, 271:21353–21358; Transformation related protein 53 (Trp53), Boehme SA, et al. J Immunol 1996, 156:4075–4078; Neuronal nitric oxide synthase (Nos1),
- 20 Burnett AL, et al. Mol Medicine 1996, 2:288–296; Eph receptor A2 (Epha2), Chen J, et al. Oncogene 1996, 12:979–988; Urokinase plasminogen activator receptor (Plaur); Dewerchin M, et al. J Clin Invest 1996, 97:870–878; Growth differentiation factor 9 (Gdf9), Dong J, et al. Nature 1996, 383:531–535; Externally regulated phosphatase (Ptpn16), Dorfman K, et al. Oncogene 1996, 13:925–931; Tenascin C (Tnc), Forsberg E,
- 25 et al. Proc Natl Acad Sci U S A 1996, 93:6594–6599; Integrin alpha 1 (Itga1), Gardner H, et al. Dev Biol 1996, 175:301–313; FBJ osteosarcoma oncogene B (Fosb), Gruda MC, et al. Oncogene 1996, 12:2177–2185; Breast cancer 1 (Brcal), Hakem R, et al. Cell 1996, 85:1009–1023; Megakaryocyte-associated tyrosine kinase (Matk), Hamaguchi I, et al. Biochem Biophys Res Commun 1996, 224:172–179; Apolipoprotein B editing
- 30 complex 1 (Apobec1), Hirano K-I, et al. J Biol Chem 1996, 271:9887–9890; Carboxyl ester lipase, Howles PN, et al. J. Biol. Chem. 1996, 271:7196–7202; Nuclear factor, erythroid derived 2, ubiquitous (Nfe2u), Kotkow KJ, Orkin SH. Proc Natl Acad Sci U S

- A 1996, 93:3514–3518; Retinoid X receptor gamma (Rxrg), Krezel W, et al. Proc Natl Acad Sci U S A 1996, 93:9010–9014; Early growth response 1 (Egr1), Lee SL, et al. Mol Cell Biol 1996, 16:4566–4572; Adrenergic receptor, alpha 2b (Adra2b), Link RE, et al. Science 1996, 273:803–805; Angiotensin receptor 1a (Agtr1a), Matsusaka T, et al. J Clin Invest 1996, 98:1867–1877; Interleukin 3 receptor, beta chain 1 (Il3rb2), Nicola NA, et al. Blood 1996, 87:2665–2674; Transformation related protein 53 (Trp53), Ohashi M, et al. Jpn J Cancer Res 1996, 87:696–701; Leukemia-associated gene (Lag), Schubart U K, et al. J Biol Chem 1996, 271:14062–14066; Glutathione peroxidase 1 (Gpx1), Spector A, et al. Exp Eye Res 1996, 62:521–540; Arachidonate 12-lipoxygenase, leukocyte (Alox12l), Sun D, Funk CD. J Biol Chem 1996, 271:24055–24062; Complement component 3 (C3), Sylvestre D, et al. J Exp Med 1996, 184:2385–2392; CD30 antigen (Cd30), Texido G, et al. Eur J Immunol 1996, 26:1966–1969; Fc receptor, IgE, low affinity II, alpha polypeptide (Fcer2a), Immunoglobulin heavy chain 5 (delta-like heavy chain) (Igh-5), Interleukin 4 (IL4) and Terminal-deoxynucleotidyl transferase, Texido G, et al. Eur J Immunol 1996, 26:1966–1969; Apolipoprotein A-II (Apoa2), Weng W, Breslow JL. Proc Natl Acad Sci U S A 1996, 93:14788–14794; Amyloid beta (A4) precursor protein (App), Zheng H, et al. Ann N Y Acad Sci 1996, 777:421–426; cAMP responsive element binding protein 1 (Creb1), Blendy JA, et al. Brain Res 1995, 681:8–14; Bradykinin receptor, beta 2 (Bdkrb2), Borkowski JA, et al. J Biol Chem 1995, 270:13706–13710; Growth factor response protein (Gfrp), Crawford PA, et al. Mol Cell Biol 1995, 15:4331–4336; Ciliary neurotrophic factor (Cntf), de Chiara TM, et al. Cell 1995, 83:313–322; Cyclin dependent kinase inhibitor 1A (P21) (Cdkn1a); Deng C, et al. Cell 1995, 82:675–684; Granzyme A (Gzma), Ebnet K, et al. EMBO J 1995, 14:4230–4239; Very low density lipoprotein receptor (Vldlr), Frykman PK, et al. Proc Natl Acad Sci U S A 1995, 92:8453–8457; Apolipoprotein E (ApoE) / Apolipoprotein A-I (Apoa1), Goodrum JF, et al. J Neurobiol 1995, 64:408–416; Nitric oxide synthase 1, neuronal (Nos1), Ichinose F, et al. Anesthesiology 1995, 83:101–108; Nitric oxide synthase 2, inducible, macrophage (Nos2), Laubach VE, et al. Proc Natl Acad Sci U S A 1995, 92:10688–10692; Peroxisome proliferator activated receptor alpha (Ppara), Lee SS-T, et al. Mol Cell Biol 1995, 15:3012–3022; Growth factor response protein (Gfrp), Lee SL, et al. Science 1995, 269:532–535; H19 fetal liver mRNA (H19) / Insulin-like growth factor 2 (Igf2), Leighton PA, et al. Nature 1995, 375:34–39; Retinoic

- acid receptor beta (RAR-beta), Luo J, et al. *Mech Dev* 1995, 53:61-71; Metallothionein 1 (Mt1) / Metallothionein 2 (Mt2), Philcox JC, et al. *Biochem J* 1995, 308:543-546; Heme oxygenase (decycling) 2 (Hmox2), Poss KD, et al. *Neuron* 1995, 15:867-873; H1-0 histone (H1fv), Sirotkin AM, et al. *Proc Natl Acad Sci U S A* 1995, 92:6434-6438;
- 5 Creatine kinase, mitochondrial 1, ubiquitous (Ckmt1), Steeghs K, et al. *Biochim Biophys Acta* 1995, 1230:130-138; Tenascin C (Tnc), Steindler DA, et al. *J Neurosci* 1995, 15:1971-1983; Ia-associated invariant chain (Ii), Takaesu NT, et al. *Immunity* 1995, 3:385-396; Neuroblastoma ras oncogene (Nras), Umanoff H, et al. *Proc Natl Acad Sci USA* 1995, 92:1709-1713; Receptor-associated protein of the synapse, 43 kDa (Rapsn),
- 10 Willnow TE, et al. *Proc Natl Acad Sci USA* 1995, 92:4537-4541; Vitronectin (Vtn), Zheng X, et al. *Proc Natl Acad Sci U S A* 1995, 92:12426-12430; Preproacrosin (Acr), Baba T, et al. *J Biol Chem* 1994, 269:31845-31849; Vimentin (Vim), Colucci GE, et al. *Cell* 1994, 79:679-694; Tumor necrosis factor receptor superfamily, member 1b (Tnfrsf1b), Erickson SL, et al. *Nature* 1994, 372:560-563; Cellular retinoic acid binding
- 15 protein I (Crabp1), Gorry P, et al. *Proc Natl Acad Sci USA* 1994, 91:9032-9036; cAMP responsive element binding protein 1 (Creb1), Hummler F, et al. *Proc Natl Acad Sci USA* 1994, 91:5647-5651; Pore forming protein (Pfp), Kagi D, et al. *Nature* 1994, 369:31-37; Src-related kinase lacking C-terminal regulatory tyrosine and N-terminal myristylation sit (Srms), Kohmura N, et al. *Mol Cell Biol* 1994, 14:6915-6925; CD3
- 20 polypeptide zeta (Cd3z) / Solute carrier family 22, member 1 (Slc22a1), Koyasu S, et al. *EMBO J* 1994, 13:784-797; Pore forming protein (Pfp), Lowin B, et al. *Proc Natl Acad Sci USA* 1994, 91:11571-11575; Retinoic acid receptor, beta (Rarb), Retinoic acid receptor beta2 (RARbeta2), Mendelsohn C, et al. *Dev Biol* 1994, 166:246-258; Transthyretin (Ttr), Palha JA, et al. *J Biol Chem* 1994, 269:33135-33139; Procollagen,
- 25 type X, alpha 1 (Col10a1), Rosati R, et al. *Nature Genet* 1994, 8:129-135; P glycoprotein 3 (Pgy3), Schinkel AH, et al. *Cell* 1994, 77:491-502; Yamaguchi sarcoma viral (v-yes) oncogene homolog (Yes), Stein PL, et al. *Genes Dev* 1994, 8:1999-2007; Fc receptor, IgE, high affinity II, alpha polypeptide (Fcer2a), Stief A, et al. *J Immunol* 1994, 152:3378-3390; Pore forming protein (Pfp), Walsh CM, et al. *Proc Natl Acad Sci*
- 30 *USA* 1994, 91:10854-10858; CD2 antigen (Cd2), Evans CF, et al. *J Immunol* 1993, 151:6259-6264; Mannose-6-phosphate receptor, cation dependent (M6pr), Koster A, et al. *EMBO J* 1993, 12:5219-5223; Retinoic acid receptor, alpha (Rara), Li E, et al. *Proc*

- Natl Acad Sci USA 1993, 90:1590–1594; Lufkin T, et al. Proc Natl Acad Sci USA 1993, 90:7225–7229; Retinoic acid receptor, gamma (Rarg), Lohnes D, et al. Cell 1993, 73:643–658; Tumor necrosis factor receptor 1 (TNF-R-1) (Tnfr1), Pfeffer K, et al. Cell 1993, 73:457–467; Lectin, galactose binding, soluble 1 (Lgals1), Poirier F, Robertson EJ. Development 1993, 119:1229–1236; Synapsin I (Syn1), Rosahl TW, et al. Cell 1993, 75:661–670; Tumor necrosis factor receptor 1 (Tnfr1), Rothe J, et al. Nature 1993, 364:798–802; Beta-2 microglobulin (B2m), Correa I, et al. Proc Natl Acad Sci USA 1992, 89:653–657; CD2 antigen (Cd2), Killeen N, et al. EMBO J 1992, 11:4329–4336; Apolipoprotein E (ApoE), Piedrahita JA, et al. Proc Natl Acad Sci USA 1992, 89:4471–4475; Myogenic differentiation 1 (Myod1), Rudnicki MA, et al. Cell 1992, 71:383–390; Tenascin C (Tnc), Saga Y, et al. Genes Dev 1992, 6:1821–1831; Beta-2 microglobulin (B2m), Sanjuan N, et al. J Virol 1992, 66:4587–4590; Neuroblastoma myc-related oncogene 1 (Nmyc1), Stanton BR, et al. Genes Dev 1992, 6:2235–2247; and Hemoglobin alpha chain complex (Hba), Popp RA, et al. Genetics 1983, 105:157–167.
- 15       Some preferred single-copy, non-essential genes with no phenotypes of the present invention include, but are not limited to, the following: Moesin (Msn), Doi Y., et al., J Biol Chem 1999, 274:2315–2321; Plasminogen activator inhibitor, type II (Planh2) and Planh1, Dougherty K.M., Proc Natl Acad Sci USA 1999, 96:686–691; Nuclear receptor coactivator 1 (Nco1), Qi C, et al. (1999) Proc Natl Acad Sci USA 96:1585–1590; Nuclear factor of kappa light chain gene enhancer in B-cells inhibitor, alpha and beta (Nfkb1a and Nfkb1b), Cheng JD, et al. (1998) J Exp Med 6:1055–1062; H19 fetal liver mRNA (H19), Jones BK, et al. (1998) Genes Dev 12:2200–2207; Prion protein (Pmp), Lipp HP, et al. Behav Brain Res 1998, 95:47–54; Centromere autoantigen B (Cenpb), Perez-Castro AV, et al. Dev Biol 1998, 201:135–143; Placentae and embryos oncofetal gene (Pem), Pitman JL, et al. Dev Biol 1998, 202:196–214; Externally regulated phosphatase (Ptpn16), Dorfman K, et al. Oncogene 1996, 13:925–931; Transformation related protein 53 (Trp53), Ohashi M, et al. Jpn J Cancer Res 1996, 87:696–701; H1-0 histone (H1fv), Sirotkin AM, et al. Proc Natl Acad Sci U S A 1995, 92:6434–6438; Creatine kinase, mitochondrial 1, ubiquitous (Ckmt1), Steeghs K, et al. Biochim Biophys Acta 1995, 1230:130–138; Neuroblastoma ras oncogene (Nras), Umanoff H, et al. Proc Natl Acad Sci USA 1995, 92:1709–1713; Vitronectin (Vtn), Zheng X, et al. Proc Natl Acad Sci U S A 1995, 92:12426–12430; Vimentin (Vim),
- 20
- 25
- 30



Colucci GE, et al. Cell 1994, 79:679–694; Cellular retinoic acid binding protein I (Crabp1), Gorry P, et al. Proc Natl Acad Sci USA 1994, 91:9032–9036; Retinoic acid receptor beta2 (RARbeta2), Mendelsohn C, et al. Dev Biol 1994, 166:246–258; Retinoic acid receptor, alpha (Rara), Li E, et al. Proc Natl Acad Sci USA 1993, 90:1590–1594,  
5 Lufkin T, et al. Proc Natl Acad Sci USA 1993, 90:7225–7229; Lectin, galactose binding, soluble 1 (Lgals1), Poirier F, Robertson EJ. Development 1993, 119:1229–1236; Myogenic differentiation 1 (Myod1), Rudnicki MA, et al. Cell 1992, 71:383–390; and Tenascin C (Tnc), Saga Y, et al. Genes Dev 1992, 6:1821–1831.

In view of the guidance of the present specification, one of ordinary skill in the  
10 art can select similar, suitable, single-copy, non-essential genes in mice and other cell types/organisms.

#### Assembly of Targeting Cassettes

The targeting cassettes described herein can be constructed utilizing  
15 methodologies known in the art of molecular biology (see, for example, Ausubel or Maniatis) in view of the teachings of the specification. As described above, the targeting constructs are assembled by inserting, into a suitable vector backbone, polynucleotides encoding a reporter, such as a light-generating protein, *e.g.*, a luciferase gene, operably linked to a promoter of interest; a sequence encoding a positive selection marker; and,  
20 optionally a sequence encoding a negative selection marker. In addition, the targeting cassette contains insertion sites such that sequences targeting a single-copy, non-essential gene can be readily inserted to flank the sequence encoding positive selection marker and luciferase-encoding sequence.

A preferred method of obtaining polynucleotides, suitable regulatory sequences  
25 (*e.g.*, promoters) is PCR. General procedures for PCR as taught in MacPherson et al., PCR: A PRACTICAL APPROACH, (IRL Press at Oxford University Press, (1991)). PCR conditions for each application reaction may be empirically determined. A number of parameters influence the success of a reaction. Among these parameters are annealing temperature and time, extension time, Mg<sup>2+</sup> and ATP concentration, pH, and the relative  
30 concentration of primers, templates and deoxyribonucleotides. Exemplary primers are described below in the Examples. After amplification, the resulting fragments can be



detected by agarose gel electrophoresis followed by visualization with ethidium bromide staining and ultraviolet illumination.

In one embodiment, PCR can be used to amplify fragments from genomic libraries. Many genomic libraries are commercially available. Alternatively, libraries  
5 can be produced by any method known in the art. Preferably, the organism(s) from which the DNA is has no discernible disease or phenotypic effects. This isolated DNA may be obtained from any cell source or body fluid (*e.g.*, ES cells, liver, kidney, blood cells, buccal cells, cerviovaginal cells, epithelial cells from urine, fetal cells, or any cells present in tissue obtained by biopsy, urine, blood, cerebrospinal fluid (CSF), and tissue  
10 exudates at the site of infection or inflammation). DNA is extracted from the cells or body fluid using known methods of cell lysis and DNA purification. The purified DNA is then introduced into a suitable expression system, for example a lambda phage.

Another method for obtaining polynucleotides, for example, short, random nucleotide sequences, is by enzymatic digestion. As described below in the Examples,  
15 short DNA sequences generated by digestion of DNA from vectors carrying genes encoding luciferase (yellow green or red).

Polynucleotides are inserted into vector genomes using methods known in the art. For example, insert and vector DNA can be contacted, under suitable conditions, with a restriction enzyme to create complementary or blunt ends on each molecule that can pair  
20 with each other and be joined with a ligase. Alternatively, synthetic nucleic acid linkers can be ligated to the termini of a polynucleotide. These synthetic linkers can contain nucleic acid sequences that correspond to a particular restriction site in the vector DNA. Other means are known and, in view of the teachings herein, can be used.

25 The final constructs can be used immediately (*e.g.*, for introduction into ES cells), or stored frozen (*e.g.*, at -20°C) until use. Preferably, the constructs are linearized prior to use, for example by digestion with suitable restriction endonucleases.

### Transgenic Animals

The targeting constructs containing the light generating protein coding sequences (e.g., luciferase genes) are introduced into a pluripotent cell (e.g., ES cell, Robertson, E. J., In: Current Communications in Molecular Biology, Capecchi, M. R. (ed.), Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1989), pp. 39-44). Suitable ES cells may be derived or isolated from any species or from any strain of a particular species. Although not required, the pluripotent cells are typically derived from the same species as the intended recipient. ES cells may be obtained from commercial sources, from International Depositories (e.g., the ATCC) or, alternatively, may be obtained as described in Robertson, E. J., *supra*. Examples of clonally-derived ES cells lines include 129/SVJ ES cells, RW-4 and C57BL/6 ES cells (Genome Systems, Inc.).

ES cells are cultured under suitable conditions, for example, as described in Ausubel et al., section 9.16, *supra*. Preferably, ES cells are cultured on stromal cells (such as STO cells (especially SNC4 STO cells) and/or primary embryonic fibroblast cells) as described by E. J. Robertson, *supra*, pp 71-112. Culture media preferably includes leukocyte inhibitory factor ("lif") (Gough, N. M. et al., Reprod. Fertil. Dev. 1:281-288 (1989); Yamamori, Y. et al., Science 246:1412-1416 (1989), which appears to help keep the ES cells from differentiating in culture. Stromal cells transformed with the gene encoding lif can also be used.

The targeting constructs are introduced into the ES cells by any method which will permit the introduced molecule to undergo recombination at its regions of homology, for example, micro-injection, calcium phosphate transformation, or electroporation (Toneguzzo, F. et al., Nucleic Acids Res. 16:5515-5532 (1988); Quillet, A. et al., J. Immunol. 141:17-20 (1988); Machy, P. et al., Proc. Natl. Acad. Sci. (U.S.A.) 85:8027-8031 (1988)). The construct to be inserted into the ES cell must first be in the linear form. Thus, if the knockout construct has been inserted into a vector as described above, linearization is accomplished by digesting the DNA with a suitable restriction endonuclease selected to cut only within the vector sequence and not within the knockout construct sequence. If the ES cells are to be electroporated to insert the construct, the ES cells and construct DNA are exposed to an electric pulse using an electroporation machine and following the manufacturer's guidelines for use. After electroporation, the ES cells are typically allowed to recover under suitable incubation conditions. The cells

are then cultured under conventional conditions, as are known in the art, and screened for the presence of the construct.

Screening and selection of those cells into which the targeting construct has been integrated can be achieved using the positive selection marker and/or the negative  
5 selection marker in the construct. In preferred embodiments, the construct contains both positive and negative selection markers. In one aspect, methods which rely on expression of the selection marker are used, for example, by adding the appropriate substrate to select only those cells which express the product of the positive selection marker or to eliminate those cells expressing the negative selection marker. For  
10 example, where the positive selection marker encodes neomycin resistance, G418 is added to the transformed ES cell culture media at increasing dosages. Similarly, where the negative selection marker is used, a suitable substrate (*e.g.*, gancyclovir if the negative selection marker encodes HSV-TK) is added to the cell culture. Either before or after selection using the appropriate substrate, the presence of the positive and/or  
15 negative selection markers in a recipient cell can also be determined by others methods, for example, hybridization, detection of radiolabelled nucleotides, PCR and the like. In preferred embodiments, cells having integrated targeting constructs are first selected by adding the appropriate substrate for the positive and/or negative selection markers. Cells that survive the selection process are then screened by other methods, such as PCR or  
20 Southern blotting, for the presence of integrated sequences.

After suitable ES cells containing the construct in the proper location have been identified, the cells can be inserted into an embryo, preferably a blastocyst. The blastocysts are obtained by perfusing the uterus of pregnant females. In one embodiment, the blastocysts are obtained from, for example, the FVB/N strain of mice and the ES cells  
25 are obtained from, for example, the C57BL/6 strain of mice. Suitable methods for accomplishing this are known to the skilled artisan, and are set forth by, *e.g.*, Bradley et al., (1992) *Biotechnology*, 10:534-539. Insertion into the embryo may be accomplished in a variety of ways known to the skilled artisan, however a preferred method is by microinjection. For microinjection, about 10-30 ES cells are collected into a micropipet  
30 and injected into embryos that are at the proper stage of development to permit integration of the foreign ES cell containing the construct into the developing embryo.

The suitable stage of development for the embryo used for insertion of ES cells is species dependent, in mice it is about 3.5 days.

While any embryo of the right stage of development is suitable for use, it is preferred that blastocysts are used. In addition, preferred blastocysts are male and, furthermore, preferably have genes encoding a coat color that is different from that encoded by the genes ES cells. In this way, the offspring can be screened easily for the presence of the knockout construct by looking for mosaic coat color (indicating that the ES cell was incorporated into the developing embryo). Thus, for example, if the ES cell line carries the genes for black fur, the blastocyst selected will carry genes for white or brown fur.

After the ES cell has been introduced into the blastocyst, the blastocyst is typically implanted into the uterus of a pseudopregnant foster mother for gestation. Pseudopregnant females are prepared by mating with vasectomized males of the same species and successful implantation usually must occur within about 2-3 days of mating.

Offspring are screened initially for mosaic coat color where the coat color selection strategy has been employed. Southern blots and/or PCR may also be used to determine the presence of the sequences of interest. Mosaic (chimeric) offspring are then bred to each other to generate homozygous animals. Homozygotes and heterozygotes may be identified by Southern blotting of equivalent amounts of genomic DNA from mice that are the product of this cross, as well as mice that are known heterozygotes and wild type mice. Alternatively, Northern blots can be used to probe the mRNA to identify the presence or absence of transcripts encoding either the replaced gene, the light generating protein coding sequence (e.g., luciferase gene), or both. In addition, Western blots can be used to assess the level of expression of the luciferase protein with an antibody against the luciferase gene product. Finally, in situ analysis (such as fixing the cells and labeling with antibody) and/or FACS (fluorescence activated cell sorting) analysis of various cells from the offspring can be conducted using suitable (e.g., anti-luciferase) antibodies to look for the presence or absence of the targeting construct.

In one embodiment of the present invention, the animals are from the C57BL/6 mouse strain. This strain develops a variety of tumors and has been used to develop a number of tumor cells lines, for example, B16 melanoma cells (including, B16F10, B16D5, and B16F1), Lewis lung carcinoma cells (including, LLC, LLC-h59), T241

mouse fibrosarcoma cells, RM-1 and pTC2 mouse prostate cancer cells, and MCA207 mouse sarcoma cells. These cell lines have been extensively used for *in vivo* tumor biology studies after injection into C57BL/6 mice. The generated targeted transgenic mice in the Examples are in C57BL/6 genetic background and these animals are suitable  
5 for injection or implantation of such tumor cells, as well as other tumor cells described in literature that are immunocompatent for C57BL/6 mice. Thus, the transgenic animals can then be used, for example, to monitor, *in vivo*, tumor progression (*e.g.*, growth) and the efficacy of therapies on tumor regression. For example, where the transgenic animal is tumor-susceptible, it is monitored for expression of a reporter, *e.g.*, luciferase, which is  
10 indicative of tumorigenesis and/or angiogenesis. The monitoring of expression of luciferase reporter expression cassettes using non-invasive whole animal imaging has been described (Contag, C. et al, U.S. Patent No. 5,650,135, July 22, 1997; Contag, P., et al, *Nature Medicine* 4(2):245-247, 1998; Contag, C., et al, *OSA TOPS on Biomedical Optical Spectroscopy and Diagnostics* 3:220-224, 1996; Contag, C.H., et al,  
15 *Photochemistry and Photobiology* 66(4):523-531, 1997; Contag, C.H., et al, *Molecular Microbiology* 18(4):593-603, 1995). Such imaging typically uses at least one photo detector device element, for example, a charge-coupled device (CCD) camera.

The transgenic animals described herein can also be used to determine the effect of an analyte (*e.g.*, therapy), for example on tumor progression where the promoter  
20 induces light generating protein (*e.g.*, luciferase) expression when a tumor develops. Methods of administration of the analyte include, but are not limited to, injection (subcutaneously, epidermally, intradermally), intramucosal (such as nasal, rectal and vaginal), intraperitoneal, intravenous, oral or intramuscular. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal  
25 applications. Dosage treatment may be a single dose schedule or a multiple dose schedule. For example, the analyte of interest can be administered over a range of concentration to determine a dose/response curve. The analyte may be administered to a series of test animals or to a single test animal (given that response to the analyte can be cleared from the transgenic animal).

30 The following examples are intended only to illustrate the present invention and should in no way be construed as limiting the subject invention.

## EXAMPLES

### **Example 1**

#### **Generating the Targeting Cassette and Vector**

5

##### **A. Creation of the Backbone Vector**

**pTK53:** The 0.5 kb mouse phosphoglycerate kinase 1 promoter was amplified with PGK primers (PGKF, SEQ ID NO:1:

ATCGAATTCTACCGGGTAGGGGAGGCGCTTT; PGKR, SEQ ID NO:2:

10 GGCTGCAGGTCGAAAGGCCCGGAGATGAGG) using mouse genomic DNA (Genome Systems, Inc., St. Louis, MO) as template. This fragment was then double digested with EcoRI and PstI and cloned into the pKS vector (Stratagene, La Jolla, California) which was linearized with the same enzymes. The neomycin gene was amplified with NeoF (SEQ ID NO:3:

15 ACCTGCAGCCAATATGGGATCGGCCATTGAAC) and NeoR (SEQ ID NO:4: GGATCCGCGGCCGCCCCCAGCTGGTTCTTCCGCCTC) primers using pNTKV1907 (Stratagene) as a template. The 1.1 kb PCR fragment was double digested with PstI and BamHI and cloned into the pKS-PGK vector which was linearized with the same enzymes. This pKS-PGK-Neo vector was used to clone thymidine kinase gene as follows. Primers TKF (SEQ ID NO: 5:

GGATCCTCTAGAGTCGAGCAGTGTGGTTTT) and TKR (SEQ ID NO:6:

GAGCTCCCGTAGTCAGGTTTAGTTCGTCCG) were used to amplify the TK gene from pNTKV1907 (Stratagene). The amplified 2kb fragment was then digested with

BamHI and SacI and cloned into pKS-PGK-Neo vector that was linearized with the same

25 enzymes. This constructed vector was designated as pTK. A synthetic linker F5R5 was made after annealing of two primers (forward primer, F5R51, SEQ ID NO: 7:

GTACATTTAAATCCTGCAGG; reverse primer, F5R52, SEQ ID NO:8:

AGCTCCTGCAGGATTTAAAT). This linker was inserted between Asp718I and

HindIII sites of pTK and the new construct was designated pTK5. A second synthetic

30 linker F3R3 was made by annealing of two primers (forward primer, F3R31, SEQ ID NO:9:

GGCCCGGGCTTAATTAATGCATCATATGGTACCGTTTAAACGCGGCCGCAAG

CTTGTCGACGGCGCGCCGGCCGGCC; reverse primer, F3R32, SEQ ID NO:10:



GATCGGCCGGCCGGCGCGCCGTCGACAAGCTTGCGGCCGCGTTTAAACGGTACCATATGATGCATTAATTAAGCCCG). ). This linker was inserted between NotI and BamHI sites of pTK and the new construct was designated pTK53. Schematics of the vectors are shown in Figure 1.

5

#### **B. Introduction of Luciferase**

**pTK-LucYG and pTK-LucR:** The yellow green luciferase gene was isolated from pGL3 vector (Promega) as a HindIII-SalI fragment and was cloned into pGK53 that was linearized with the same enzymes. The new construct was designated pTK-LucYG  
10 (8931 bp), shown in Figure 2.

The red luciferase gene was isolated from pGL3-red vector (Dr. Christopher Contag, Stanford University, Stanford, Calif.) as a HindIII-SalI fragment and was cloned into pGK53 that was linearized with the same enzymes. The new construct was designated pTK-LucR (8931 bp), shown in Figure 2.

15

### **Example 2**

#### **Insertion of Targeting Sequences**

**A. Generation of vitronectin targeting vector:** The targeting construct  
20 pTKLR-Vn was generated by inserting vitronectin (VN) DNA sequences into pTK-LucR vector.

Vitronectin (VN) is an abundant glycoprotein present in plasma and the extracellular matrix of most tissues. In a previous study, it was shown that heterozygous mice carrying one normal and one null VN allele and homozygous null mice completely  
25 deficient in vitronectin demonstrate normal development, fertility, and survival. This suggests that VN is not essential for cell adhesion and migration during normal mouse development (Zheng, X., et al., Proc Natl Acad Sci U S A 1995 92:12426-30). Mouse vitronectin genomic DNA sequence of 5004 bp was obtained from GenBank database (Accession number X72091). Based on this sequence, a 1.63 kb 3'end vitronectin  
30 fragment was amplified (reverse primer, VN1R, SEQ ID NO:11: CTGTATTAAATCTGCCCACCCTATTCAGGACAGTAGTC; forward primer, VN1F, SEQ ID NO:12: CCAATGCATCAACCCAGCCAGGAGGAGTGCG) using mouse C57BL/6 genomic DNA as template (Genome Systems, Inc., St. Louis, MO).

This fragment was digested with *SwaI* and *NsiI* and cloned into pTK-LucR (linearized with *SwaI* and *SbfI*). This construct was designated as pTK-LucR3. Subsequently, a 2.35 kb 5'end vitronectin fragment was amplified (reverse primer, VN2R, SEQ ID NO:13: AACGCGTCGACTTCGGAGATGTTTCGGGGATAACCAGG; forward primer, VN2F, SEQ ID NO:14: TTGGCGCGCCCCATAGAGAAGAGACACCAAAGGCACGCTC) using mouse C57BL/6 genomic DNA as template. This fragment was digested with *Sall* and *AscI* and cloned into pTK-LucR vector that was linearized with *Sall* and *AscI*. This construct was designated as pTKLR-Vn. Figure 3 shows the restriction map of pTKLR-Vn vector. The polylinker between the neomycin gene and red luciferase gene is used to insert the VEGF promoter or other promoters of interests. The predicted homologous recombination between pTKLR-Vn and vitronectin gene is illustrated in Figure 3A. Upon insertion of the VEGF-LucR transgene cassette, the endogenous vitronectin gene is destroyed. Figure 3B shows the genomic DNA sequence of VN.

15

**B. Generation of Fos targeting vector:** The targeting construct pTKLG-Fos was generated by inserting FosB DNA sequences into pTK-LucYG vector.

FosB is one of the members of the Fos family. It plays a functional role in transcriptional regulation. It has been shown that FosB mice are born at a normal frequency, are fertile and present no obvious phenotypic or histologic abnormalities (Gruda et al (1996) *Oncogene* 12:2177-2185). A 28.8 kb genomic region that contains mouse FosB DNA sequence was obtained from GenBank database (Accession number AF093624).

Using this sequence, a 1.71 kb 5'end FosB fragment was amplified (forward primer, FosB1F, SEQ ID NO:15: CTGTATTTAAATCCCGTTTCTCACTGTGCCTGTGTC; reverse primer, FosB1R, SEQ ID NO:16: GTCTCCTGCAGGCTTCCTCCTCCTTGTTTCCTTGCG) using mouse C57BL/6 genomic DNA as template. This fragment was digested with *SwaI* and *SbfI* and cloned into pTK-LucYG vector that was linearized with *SwaI* and *SbfI*. This construct was designated as pTK-LucYG3. Subsequently, a 1.58 kb 3'end FosB fragment was amplified (forward primer, FosB2F, SEQ ID NO:17: AACGCGTCGACGGATGGGATTGACCCCCAGCCCTC; reverse primer, FosB2R,

30

SEQ ID NO:18: TTGGCGCGCCCCTTGCCTCCACCTCTCAAATGC) using mouse C57BL/6 genomic DNA as template. This fragment was digested with *Sall* and *AscI* and cloned into pTK-LucYG vector that was linearized with *Sall* and *AscI*. This construct was designated as pTKLG-Fos (Figure 4). The polylinker between the neomycin gene and red luciferase gene is used to insert the VEGFR2 promoter (Example 3, Figure 5A), Tie2 promoter (Example 3, Figure 5B), as well as, other promoters of interests. The predicted homologous recombination between the targeting vector bearing the VEGFR2 promoter (Figure 5A) or the Tie2 promoter (Figure 5B) and FosB gene is also illustrated. As shown in the Figures, the VEGFR2-LucYG transgene cassette and Tie2-LucYG transgene cassette is inserted downstream of FosB gene translational stop signal. Therefore, the targeted transgenic mice should still have a functional FosB gene while expressing the transgenes. Figure 4B shows the DNA sequence of FosB.

### Example 3

#### 15 Insertion of Promoter Sequences of Interest

A. pTKLR-Vn/VEGF: Mouse VEGF genomic DNA sequence of 2240 bp that contains a partial VEGF promoter region was obtained from GenBank (accession number: U41383). Accordingly, primers were designed to amplify a 0.69 kb (VF1-VR1A; Table 1) and a 0.98 kb fragment (VF2-VR2; Table 1). It was confirmed that each pair of primers can amplify the predicted product using mouse 129SvJ genomic DNA as template.

Table 1

Name	Seq Id No	Sequence
VF1	19	ACCTC ACTCT CCTGT CTCCC CTGAT TCCCA A
VR1A	20	GCTCT GGCGG TCACC CCAA AAGCA
VF2	21	CCCTT TCCAA GACCC GTGCC ATTTG AGC
VR2	22	ACTTT GCCCC TGTCC CTCTC TCTGT TCGC
KF1	23	GCTGC GTCCA GATTG GCTCT CAGAT GCG
KR1	24	TTCTC AGGCA CAGAC TCCTT CTCCG TCCCT
KF2	25	CAGAT GGACG AGAAA ACAGT AGAGG CGTTG GC
KR2	26	GAGGA CTCAG GGCAG AAAGA GAGCG
TF3	27	AGCTT AGCCT GCAAG GGTGG TCCTC ATCG
TF2	28	CAAAT GCACC CCAGA GAACA GCTTA GCCTG C
TR1	29	GCTTT CAACA ACTCA CAACT TTGCG ACTTC CCG

Conditions for PCR amplification are shown in Figure 6. These primers were used for PCR screening of mouse 129/SvJ genomic DNA BAC (bacterial artificial chromosome) library (Genome Systems, Inc., St. Louis, MO). The library, on average, contained inserts of 120 kb with sizes ranging between 50 kb to 240 kb. A large genomic DNA fragment that contained VEGF promoter region was obtained. Southern blot analysis was performed to map the VEGF promoter region. A unique HindIII restriction site was mapped approximately 7.8 kb upstream of the ATG translational start codon of the VEGF gene. The sequences between HindIII and ATG translational start codon are inserted into the polylinker of pTKLR-Vn vector to finish the construction of targeting vector that contains VEGF-LucR transgene (Figure 3A).

#### 15 B. pTKLG-Fos/VEGFR2

A mouse VEGFR2 genomic DNA sequence of 1079 bp which contains a partial VEGFR2 promoter region was published previously (Ronicke, et al., (1996) *Cir. Res.* 79:277-285). Accordingly, primers were designed to amplify a 0.45 kb (KF1-KR1A; Table 1) and a 0.58 kb fragment (KF2-KR2; Table 1). It was confirmed that each pair of primers can amplify the predicted product using mouse 129SvJ genomic DNA as

template. DNA sequences for these primers are shown in Table 1 above and PCR amplification conditions are shown in Figure 6. These primers were used for PCR screening of mouse 129/SvJ genomic DNA BAC library. From the screening, a BAC clone that contained a large genomic DNA fragment was obtained. Based on the published restriction map of VEGFR2, a 4.5 kb HindIII-XbaI fragment that covers the VEGFR2 promoter region was subcloned from the VEGFR2 BAC clone into the pBluescriptSK vector (Stratagene, La Jolla, CA) that was linearized with HindIII and XbaI. The VEGFR2 promoter sequences of 4.1 kb, spanning from a HindIII site to the ATG translational start codon, are subcloned into the polylinker of pTKLG-Fos vector to construct the targeting vector that contains VEGFR2-LucYG transgene (Figure 5A).

### C. pTKLG-Fos/Tie2

Mouse Tie2 genomic DNA sequence of 477 bp which contain a partial Tie2 promoter region has been published previously (Fadel et al (1998) *Biochem. J.* 330(Pt. 1):335-343). Accordingly, primers were designed to amplify a 0.45 kb (TF3-TR1; Table 1) and a 0.47 kb fragment (TF2-TR1; Table 1). It was confirmed that each pair of primers amplified the predicted product using mouse 129SvJ genomic DNA as template. DNA sequences for these primers are shown in Table 1 above. PCR amplification conditions are shown in Figure 6. These primers are used for PCR screening of mouse 129SvJ genomic DNA BAC library. From the screening, a BAC clone containing a large genomic DNA fragment of the Tie2 promoter region was obtained. Based on the published Tie2 genomic DNA restriction map (see, Dumont et al., *supra*), a 10.5 kb Asp718-EcoRV fragment containing the Tie2 promoter region was subcloned from the Tie2 BAC clone into the pSK vector that was linearized with Asp718 and EcoRV. The Tie2 promoter sequences of about 6.8 kb, spanning from the Asp718 site to the ATG translational start codon is subcloned into the polylinker of pTKLG-Fos vector to construct the Tie2-LucYG targeting vector (Figure 5B).

**Example 4****Generation of Transgenic Mice Carrying the Constructs of the Present Invention**

5       **A. General Procedure:** Figure 7 depicts a generalized description of generation of transgenic mice using the targeted transgenic vectors described in Example 3. Details regarding embryonic stem (ES) cell culture, transfection, blastocyst injection and implantation to a pseudopregnant foster are described, for example, in Hogan et al (1994) "Manipulating the Mouse Embryo, A Laboratory Manual. Second Edition", Cold Spring  
10   Harbour Laboratory Press.

After construction the targeted transgenic construct are transfected into C57BL/6 embryonic stem (ES) cells. (Genome System Inc., Genome Systems, Inc., St. Louis, MO) through electroporation. The antibiotic G418 is used to select for cells in which the DNA construct containing the Neo gene is integrated, either randomly or by homologous  
15   recombination. The nucleoside analog gancyclovir is converted by TK to a cytotoxic derivative. DNA that has integrated by homologous recombination lose the TK gene and are resistant to the drug, whereas cells that have incorporated the DNA randomly are likely to retain the TK gene. Thus, cells containing random integrations into a chromosomal location that allows the expression of the TK gene are killed. The G418  
20   and gancyclovir resistant clones are then be screened by PCR and Southern blot analysis and those that have homologous DNA recombination is used for FVB/N blastocyst injection (Genome System, Inc.). Between 4-16 blastocysts are transferred to the uterus of a pseudopregnant foster mother. The pups are typically born 17 days after the transfer. Either random bred mice or F1 hybrid mice make suitable recipients. Females of certain  
25   random-bred stocks (*e.g.*, CD1 mice, from Charles River Laboratories) have very large ampullae, which makes oviduct transfer easier. These mice also generally make good mothers. Alternatively, F1 hybrid females (*e.g.*, B6 x CBA F1) can be used as recipients. Although their ampullae are smaller, make exceptionally good mothers, rearing litters as small as two pups. See, for example, Hogan et al. (1994), *supra*.



**B. Screening for homologous DNA recombination positive ES cells**

- 1) **pTKLG-Fos/VEGFR2:** Analysis of homologous DNA recombination between pTKLG-Fos/VEGFR2 targeting vector and the FosB gene is carried out using Southern blot analysis as shown in Figure 8. Genomic DNA prepared from G418 resistant ES  
5 cells is digested with PvuII and probed with probe A to confirm the 5'end DNA recombination. PvuII digestion of DNA bearing homologous recombination reveals two separate bands of 8.2 and 4.0 kb, whereas digestion of DNA from homologous recombination negative clones reveals only the 8.2 kb band. The 3'end of DNA recombination is tested by hybridizing NotI digested DNA with probe B. NotI digestion  
10 of DNA bearing homologous recombination will reveal two separate bands of >8.2kb and 5.0 kb, whereas digestion of DNA from homologous recombination negative clones will only reveal the >8.2 kb band. Once homologous DNA recombination is confirmed, positive clones is selected for FVB/N blastocyst injection.
- 2) **pTKLG-Fos/Tie2:** Analysis of homologous DNA recombination between pTKLG-Fos/Tie2 targeting vector and the FosB gene is analyzed by Southern blot in a similar manner as described above for pTKLG-Fos/VEGFR2. Once homologous DNA recombination is confirmed, positive clones are selected for FVB/N blastocyst injection.
- 3) **PTKLR-Vn/VEGF:** Analysis of homologous DNA recombination between pTKLR-Vn/VEGF targeting vector and the vitronectin gene is analyzed by PCR. DNA  
20 primers designed according to the predicted homologous recombination, are listed in Table 2.

**Table 2**  
PCR primers for analysis of homologous DNA recombination between pTKLR-Vn/VEGF targeting vector and the vitronectin gene

5	<b>5'end primers</b>
	<b>F51</b> 5'- CCCAGTGTCTCTGATTTAGGGAGAGCACCTGAG -3' SEQ ID NO:30
	<b>R51</b> 5'- CCAGACTGCCTTGGGAAAAGCGCCTC -3' SEQ ID NO:31
	<b>F52</b> 5'- CAGTGAGAGTCTTCTCTGTCCCTCAATCGGTTCTG -3' SEQ ID NO:32
	<b>R52</b> 5'- TGGATGTGGAATGTGTGCGAGGCCAG -3' SEQ ID NO:33
	<b>3'end primers</b>
	<b>F31</b> 5'- AATCAAAGAGGCGAACTGTGTGTGAGAGGTCC -3' SEQ ID NO:34
	<b>R31</b> 5'- CGGCTCCCCAAAATGTGGAAGCAAGC -3' SEQ ID NO:35
	<b>F32</b> 5'- GAATCCATCTTGCTCCAACACCCCAACATC -3' SEQ ID NO:36
	<b>R32</b> 5'- CGCCTCCTCTCCCCAGTCTCCCCTTG -3' SEQ ID NO:37

Primers F51-R51 and F52-R52 amplify a 1799 bp and a 1841 bp DNA fragment respectively from the 5'end of the transgene that is integrated into the vitronectin site through homologous DNA recombination, whereas primers F31-R31 and F32-R32  
10 amplify a 3549 bp and a 3428 bp DNA fragment respectively from the 3'end of the transgene that is integrated into the vitronectin site through homologous DNA recombination. Clones that allow successful amplification of both the 5'end and 3'end of the integrated transgene are selected for FVB/N blastocyst injection.

15 **C. Analysis of chimeric mice**

The pups developed from injected blastocysts contain chimeras, as can be identified by their agouti coat color when an ES cell derived from a mouse having a dark coat color (e.g., C57BL/6) is injected into the blastocyst of a light coat color animal (e.g., FVB/N, genotype B/B). DNA analysis (e.g., Southern blotting, PCR) is conducted to  
20 further confirm the presence of the transgene in these pups as described above in Section B. These animals may be obtained commercially, for example from The Jackson Laboratory, Bar Harbor, MN.

#### **D. Generating targeted transgenic C57BL/6 mice with white coat color**

Breeding of the chimeric mice generates homozygous targeted transgenic mice, as depicted in Figure 9. The targeted mice are used to monitor gene expression through the measurement of luciferase mediated light emission from the mice. In a preferred embodiment, the targeted mouse has a light coat color (e.g., white coat color), because the black colored coat (an example of a dark coat color) of C57BL/6 mice can absorb light emitted from the body and may interfere the sensitivity of the bioluminescence assay. An inbred mouse strain C57BL/6-Tyr C2j/+ strain (Jackson Laboratory, Bar Harbor, MN) is available for this purpose. This strain of mice have white color coat, yet they still have the same genetic background as C57BL/6 mice except that the gene responsible for the black coat color is mutated. Unfortunately, C57BL/6-Tyr C2j/+ ES cells are not currently available. Therefore, the designed breeding program illustrated in Figure 9 is aimed to generate mice that are homozygous for the target transgene and have white coat color. C57BL/6 ES cells are prepared as described above and introduced into a suitable blastocyst (e.g., from the FVB/N strain of mice). The blastocysts are implanted into a foster mother. Chimeric mice are shown in Figure 9 as white animals with black and green patches. Chimeric animals are bred with C57BL/6-Tyr C2j/+ mice to create F1 hybrids. Subsequent breeding of the F1 hybrids generates several type of mice, including the one that is homozygous for the target transgene and has a white coat color (shown in Figure 9 as b/b; L/L), which is used for *in vivo* gene regulation monitoring.

A C57BL/6 mouse and a C57BL/6-Tyr C2j/+ mouse are considered to be substantially isogenic. Accordingly, the method of the present invention exemplified in Figure 9 provides a means for generating breeding groups of substantially isogenic mice in a selected genetic background carrying at least one transgene of interest.

#### **E. Dual luciferase targeted transgenic mice**

As described above, two targeting vectors are generated. PTKLR-Vn carries a red luciferase gene and is targeted into vitronectin locus. PTKLG-Fos carries a yellow-green luciferase gene and is targeted into FosB locus. A number of promoters, including VEGF promoter, VEGFR2 promoter, and Tie2 promoter are cloned into these vectors, as described above. Subsequently three type of targeted transgenic mice are generated.

VEGF mice carry VEGF promoter-red luciferase transgene (VEGF-LucR) integrated into vitronectin locus. VEGFR2 mice carry VEGFR2 promoter-yellow-green luciferase (VEGFR2-LucYG) transgene integrated into FosB locus. Tie2 mice carry Tie2 promoter-yellow-green luciferase (Tie2-LucYG) transgene integrated into FosB locus. Through a breeding program illustrated in Figure 10, dual luciferase targeted transgenic mice are produced, carrying both of the VEGF-LucR and the VEGFR2-LucYG transgenes. The degradation of luciferin by yellow-green luciferase and red luciferase generates lights that emit at 540 nM and 610 nM respectively. These wavelengths of light are measured individually using a photo-counting camera (intensified CCD). Therefore, both VEGF expression and VEGFR2 expression, for example, can then be monitored in the same mouse at the same time.

As is apparent to one of skill in the art, various modification and variations of the above embodiments can be made without departing from the spirit and scope of this invention. These modifications and variations are within the scope of this invention.

**CLAIMS**

**What is claimed is:**

- 1. A vector for use in generating a transgenic non-human mammal, said mammal having at least one single-copy, non-essential gene in its genome, said vector comprising**
  - (a) a first selectable marker and a reporter expression cassette, said reporter expression cassette comprising a transcriptional promoter element operably linked to a light-generating protein coding sequence, and**
  - (b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in said mammal's genome, said targeting polynucleotide sequences flanking (a), wherein (i) the length of the targeting polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) said transcriptional promoter element is heterologous to the single-copy, non-essential gene.**
- 2. The vector of claim 1, wherein said first selectable marker provides a positive selection.**
- 3. The vector of claim 2, wherein said first selectable marker is selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase.**
- 4. The vector of claim 3, wherein the first selectable marker is neomycin phosphotransferase II.**
- 5. The vector of any of claims 1-4, further comprising a second selectable marker, wherein at least one target polynucleotide sequence is located between said second selectable marker and (a).**
- 6. The vector of claim 5, wherein said second selectable marker provides a negative selection.**

7. The vector of claim 6, wherein said second selectable marker is selected from the group consisting of adenosine deaminase, thymidine kinase, and dihydrofolate reductase.

8. The vector of any of claims 1-7, wherein the transcriptional promoter element is selected from the group consisting of an inducible promoter, a repressible promoter, and a constitutive promoter.

9. The vector of claim 8, wherein the transcriptional promoter element is selected from the group consisting of VEGF, VEGFR, and TIE2.

10. The vector of any of claims 1-9, wherein the sequences encoding the light-generating protein are obtained from either procaryotic or eucaryotic sources.

11. The vector of claim 10, wherein the light generating protein is a luciferase.

12. The vector of any of claims 1-11, wherein said reporter expression cassette further comprises other control elements.

13. The vector of claim 12, wherein said control elements are selected from the group consisting of transcription enhancer elements, transcription termination signals, polyadenylation sequences, sequences for optimization of initiation of translation, and translation termination sequences.

14. The vector of any of claims 1-11, wherein said vector is circular.

15. The vector of claim 14, wherein said vector contains at least one restriction site whose cleavage results in a linear vector having the following arrangement of elements: targeting polynucleotide sequence - (a) - targeting polynucleotide sequences.

16. The vector of claim 5, wherein said vector is circular.



17. The vector of claim 16, wherein said vector contains at least one restriction site whose cleavage results in a linear vector having the following arrangement of elements: target polynucleotide sequence - (a) - targeting polynucleotide sequences - (second selectable marker).

18. The vector of any of claims 1-17, wherein the coding sequences of the reporter expression cassette comprise codons that are optimal for expression in a host system into which the expression cassette is to be introduced.

19. The vector of claim 18, wherein said mammal is a rodent.

20. The vector of claim 19, wherein said mammal is a mouse.

21. The vector of any of claims 1-20, wherein said targeting polynucleotide sequences from single-copy, non-essential genes are selected from the group consisting of vitronectin, *fosB*, and galactin 3.

22. A method of producing a transgenic, non-human mammal, said mammal having at least one single-copy, non-essential gene in its genome, comprising  
transfecting an embryonic stem cell of said mammal with a linear vector comprising

(a) a first selectable marker and a reporter expression cassette, said reporter expression cassette comprising a transcriptional promoter element operably linked to a light-generating protein coding sequence, and

(b) targeting polynucleotide sequences homologous to a single-copy, non-essential gene in said mammal's genome, said targeting polynucleotide sequences flanking (a), wherein (i) the length of the polynucleotide sequences are sufficient to facilitate homologous recombination between the vector and the single-copy, non-essential gene, and (ii) said transcriptional promoter element is heterologous to the single-copy, non-essential gene;

selecting embryonic stem cells which each have said first selectable marker and reporter expression cassette integrated into its genome;  
injecting said embryonic stem cells into a host embryo,  
implanting said embryo in a foster mother,  
maintaining said foster mother under conditions which allow production of an offspring that is a transgenic, non-human mammal carrying said reporter expression cassette.

23. The method of claim 22, wherein said offspring is capable of germline transmission of said reporter expression cassette.

24. The method of claim 23 further comprising breeding said offspring with a mammal, wherein the mammal is substantially isogenic with the embryonic stem cells, wherein said breeding yields transgenic F1 offspring carrying said reporter cassette.

25. The method of claim 24, further comprising breeding a first F1 offspring carrying said reporter cassette with a second F1 offspring carrying said reporter cassette, wherein said breeding yields transgenic F2 offspring carrying said reporter cassette.

26. The method of any of claims 22-25, wherein said mammal is a mouse.

27. The method of claim 26, wherein said embryonic stems cells are derived from a mouse having a dark coat color.

28. The method of claim 27, wherein said mammal substantially isogenic with the embryonic stem cells has a light coat color.

29. The method of claim 28, wherein said F2 offspring carrying said reporter cassette has a light coat color.

30. The method of claim 29, wherein said embryonic stems cells are derived from a C57BL/6 mouse having a dark coat color, and said mammal substantially

isogenic with the embryonic stem cells is a C57BL/6-Tyr C2j/+ mouse having a light coat color.

31. The method of any of claims 22-30, wherein the light generating protein is a luciferase.

32. A transgenic, non-human mammal, comprising at least one single-copy, non-essential gene in its genome, wherein (i) at least a portion of at least one single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the gene, such that the gene cannot produce a functional gene product, and (ii) said polynucleotide sequences comprise a first expression cassette which has been introduced into said mammal or an ancestor of said mammal, at an embryonic stage, said first expression cassette comprising

a first selectable marker,  
a first transcriptional promoter element heterologous to the gene, and  
light-generating protein coding sequences, wherein said light-generating protein coding sequences are operably linked to said promoter element.

33. The transgenic, non-human mammal of claim 32, wherein the single-copy, non-essential gene is selected from the group consisting of vitronectin, *fosB*, and galactin 3.

34. The transgenic, non-human mammal of any of claims 32-33, wherein the first selectable marker is selected from the group consisting of neomycin phosphotransferase II, xanthine-guanine phosphoribosyltransferase, hygromycin-B-phosphotransferase, chloramphenicol acetyltransferase, and adeninephosphoribosyl transferase.

35. The transgenic, non-human mammal of any of claims 32-34, wherein the first transcriptional promoter element is selected from the group consisting of an inducible promoter, a repressible promoter, and a constitutive promoter.

36. The transgenic, non-human mammal of claim 35, wherein the first transcriptional promoter element is selected from the group consisting of VEGF, VEGFR, and TIE2.

37. The transgenic, non-human mammal of any of claims 32-36, wherein said expression cassette further comprises other control elements.

38. The transgenic, non-human mammal of claim 37, wherein said control elements are selected from the group consisting of transcription enhancer elements, transcription termination signals, polyadenylation sequences, sequences for optimization of initiation of translation, and translation termination sequences.

39. The transgenic, non-human mammal of any of claims 32-38, wherein said non-human mammal comprises a second single-copy, non-essential gene in its genome, wherein (i) at least a portion of the second single-copy, non-essential gene is replaced by polynucleotide sequences heterologous to the second gene, such that the second gene cannot produce a functional gene product, and (ii) said polynucleotide sequences comprise a second expression cassette which has been introduced into said mammal or an ancestor of said mammal, at an embryonic stage, said first expression cassette comprising

a second selectable marker,  
a second transcriptional promoter element heterologous to the second gene, and  
light-generating protein coding sequences, wherein said light-generating protein coding sequences are operably linked to said second promoter element.

40. The transgenic, non-human mammal of claim 39, wherein the first transcriptional promoter element in the first expression cassette is different from the second transcriptional promoter element in the second expression cassette.

41. The transgenic, non-human mammal of claim 40, wherein the light-generating protein in the first expression cassette can produce a different color of light relative to the light-generating protein in the second expression cassette.

42. The transgenic, non-human mammal of any of claims 32-41, said mammal being a rodent.

43. The transgenic, non-human mammal of 42, said rodent being a mouse.

44. The transgenic, non-human mammal of any of claims 32-43, wherein the sequences encoding the light-generating protein are obtained from either procaryotic or eucaryotic sources.

45. The transgenic, non-human mammal of claim 44 wherein the light-generating protein is a luciferase.




FIG. 1






FIG. 2




FIG. 3A




FIG. 3B

1 TCAACCAAC TGTTCCTAC GTCCGCCGCC TTCCCTATTA ACTTCATGCT TAAGCAAGCC TCAACGCCGC AGGCTGTGCT CACACAGAG GAGGCTGCT  
AGGTCGGTCC ACNAGAGTGG CAGGCCGCCGCC AAGGATCAAT TGAAGTACCA ATTCCTTCCG AGTCCGCCGC TCCACACCA GGTGTCTTTC CTTCGCCGA

101 CACCAAGGCC CCAATGTCTT CTGATTTAGG GAGGACACT GAGGCCAGTG AGAGTCTTCT CTGTCCCTCA ATCCGTTCTG AATTCGCCCA CTTCGCCCTC  
GGGTGTTCCG GGTTCACAGA GACTAAATCC CTCTGCTGGA CTGCGTCCAC TCTCAGAGA GACAGGAGT TACCCAGAC TTAAAGGCTT GAGCGGAGG

201 TATCCAGGG GACAGGCTG CCAACCTAT TCAAGACGT AGTCTAAAC TGGTACCAAA CAGACTTTT ATTCCGCTGG GAGAAAGAGA TGAAGCTGCT  
AATAGGTGCC CTGTCCGAC GGTTCGATA AGTCTGTCA TCAGATTTC AGCATCGTT GTCTGAAAA TACCCGACC CTCTTCTCT ACTCCGAGGA

301 GAGGCTCAG CCACTGCTCT CTGATTCCTA CTCTCAGAG GTCCGCCAGC CCAACCAATA CTGAGCAAT GAGGCTGCT AGGAGGCTT CACAGATCC  
CTTCAGTGG GTCAACCGA GACTAAGCA GAGAGTCTC CAGGCCGTC GGTGCTTAT GACTGCTAC CTCCACCCA TCCCTCTGA GTGTCTCAG

401 ACTCCGCCGC TTCTAAGCTT GACTGCTGAG TATTTCTCT AAAGAAAGAA TGAAGAAAG GTATGTGAG ATTCTGCTG ATCTGTGCA CTGTGCCAA  
TGAGGCCGC AAGATTCCA CTGAGCAATC ATAAACAGAC TTCTTTCTT AACTTTTTC CAATACACTC TACAGCCAC TACGACAGT GAGGAGGCTT

501 GAGGATAAA GCTTTTCTT CAGAGGCA AGTGAACAT CACCAAGCAG ATATGTCTAC CATCTACAG CTGTGTTGAG CAGGCCGGA CCAAGGCTG  
CTTCTATTT CCAAAAGAA GTCTTCCCTT TCACTGTGAG GTGTTCTCTC TATTACAGTG GTAGATGTCC GACACAGTC GTGCTGCTT GGTTCGTGAC

601 CAGGCAAGCC CTAGCAAAA CCACTCTAG GAGTGAAG GGGTCCAC CTCCAGAGAA GAAAGAGAG CTCTGAATG GCTGCAAGT GCAAGTACA  
GTGCTTCCG GATCGTTTT GTTCAGATTC CTATCTTTC GGGTCCAG GTGCTCTT CTTEATCTCC GAGACTTAC GAGGCTCCA GGTGCTATG

701 AGGCGTCCA TATCATATC ATAGTTGTT TGGTTCTCA GGCCTCTC CTGCTGAG AGCAAGAGA ACCAGATTG AGGTGATGA GAGCGGAGT  
TGGTTCAGT ATAGTATAG TATCAACAC ATCCAGGAT GGTGTGAG GAGGAGCTC TTGTTCTCT TGTCTAAT TCCACTACT GCTGCCCTCA

801 TGAAGTCTG GCTGCTCTG TGGCCAGCC CTGCGCTGA AGGAGAGCC TTTCGCTTC TACCTTAGA CTCTCTTTT TTGCTTGG CAGAGTCCA  
AGCTGAGAC GAGGACAGC AGGCTGCGG GAGGCGACT TGTATGCG AAAGCGAG ATCGGATCT GAGACAAA AACCGAAC GTCTCAAGCT

901 TAAGGAGCA GTGAGTGA TCCGCCGCC CATAGCAGG TCCCTTTC CTGCGACAC ATCCAGTTC GGGTCTGA ATTCGCTTC TCTGCTCA  
ATTCTGCT CACTGCTCT AGGCGGCCG GTATGCTGC AGGTGAAGG GAGGCTGTC TACGCTCAG GGGGCTCT TACGCCAG AGAGCGGCT

1001 TCTGAGAGC GAAAGGCTT AATGCTCT GAGATCTCT GCTTGAAT CTCCAGCAG GTTGTCTC TATCTATTA TCTATCTTA TCTATCTA  
AGCATTTCT CTTGCCAA TTAGGTAAC GTTCAGAC CCAACTAAA GAGGTGCTC CACAGAGC ATAGATTAAT ACATAGAAAT ACATAGAAAT

1101 TATCTATA TCTATATC TATCTATTA TATCTACT AACTACTC CTATCTAT GTTGTCTC TATCTATTA TATCTATTA TATCTATTA  
ATAGATATA ACATAGAA TATAGATTA ATAGATTA TATAGATTA TATAGATTA TATAGATTA TATAGATTA TATAGATTA TATAGATTA

1201 OCTATTTAT TTTTCTTTG TTTCTTTGA AACAGATCT TACAGCTAC CTATGCTG TTTCAGCTC ACTATGAGC CATACTGC CTCTTACTC  
GATAAATA ACNAGAAC TTTCTTTGA TTTCTTTGA AACAGATCT TACAGCTAC CTATGCTG TTTCAGCTC ACTATGAGC CATACTGC CTCTTACTC

1301 ACAAGATCC ACTTCCCTG GTCTCTGAT CTGCGATTA AAGCATGT CCACTACAC CAGCTCCAGT AGGAGCTTA GACACATTT GCTATGCTT  
TGTTCCTAG TGAAGGCA CAGAGCTCA GCGCTAAT TTCTCTAC GGTGATGTC CAGCTCCAGT AGGAGCTTA GACACATTT GCTATGCTT

1401 GGTGAGACA CACTCTAG TCCCGAGCC CAGGCTTCC TGTCTAGAC TTTTCTCAT GGTCTCTCA CTGTCTCA CTGTCTCA CTGTCTCA  
GGATTTCT GTTTGAGTC AGGCGGCCG GGTGAGAGG ACAGATCT TTTTCTCAT GGTCTCTCA CTGTCTCA CTGTCTCA CTGTCTCA

1501 AAGGCTCAG CCGCAGGCC CTGCTCTGC TGTGTTAGC AAGTCCAG GTATGCTG CAAATGAGC CAAAGCTCA CCGCAAGG TCAACAGAG  
TTGGGAGTC GCGGCTGCC GGTCTCTGC TGTGTTAGC AAGTCCAG GTATGCTG CAAATGAGC CAAAGCTCA CCGCAAGG TCAACAGAG

1601 CAAAGTCTG CAGAGCAAA CAGCTCTGA TGGATGTT CAGATTTCA GGTGAGACT GGTGAGACT GGTGAGACT GGTGAGACT GGTGAGACT  
GTTCAGATC GTTCAGATC GTTCAGATC GTTCAGATC GTTCAGATC GTTCAGATC GTTCAGATC GTTCAGATC GTTCAGATC GTTCAGATC

1701 AAGGAGCTT TGTCTGCT CCACTACAG GATCTGCC CAGAGGCA GTTCCAAAT GTTCTCCAG CTGTGCCCT GAGCAAGCC AAGGCTCA  
TTGCTGCA AGCAGAGCA GTTCTGCT CCACTACAG GATCTGCC CAGAGGCA GTTCCAAAT GTTCTCCAG CTGTGCCCT GAGCAAGCC AAGGCTCA

1801 AAGGAGCTT TGTCTGCT CCACTACAG GATCTGCC CAGAGGCA GTTCCAAAT GTTCTCCAG CTGTGCCCT GAGCAAGCC AAGGCTCA  
TTGCTGCA AGCAGAGCA GTTCTGCT CCACTACAG GATCTGCC CAGAGGCA GTTCCAAAT GTTCTCCAG CTGTGCCCT GAGCAAGCC AAGGCTCA

1901 ATCAGGAGG ATCTGAGCA GCGAGAGAG AATCATGAA TGAAGAGAG ACTCCAGC CTGCGGCTT CTCTCCAGC CTGAGTACC TTGAGAGT  
TGTCTCTC TACTGCTT GGTCTCTC TGTCTCTC TGTCTCTC TGTCTCTC TGTCTCTC TGTCTCTC TGTCTCTC TGTCTCTC

2001 AGAGGCTTC CCGGCTCTG TACGCTGCT CAGGAGGCC GAGGCTCA TCAACATGT CTGTGCTCC ACTGAGGCT TGGAGATGT TTGCGGGA  
TCTGGAAG GCGGCTCTG TACGCTGCT CAGGAGGCC GAGGCTCA TCAACATGT CTGTGCTCC ACTGAGGCT TGGAGATGT TTGCGGGA

2101 AAGGAGCTT TGTCTGCT CCACTACAG GATCTGCC CAGAGGCA GTTCCAAAT GTTCTCCAG CTGTGCCCT GAGCAAGCC AAGGCTCA  
TTGCTGCA AGCAGAGCA GTTCTGCT CCACTACAG GATCTGCC CAGAGGCA GTTCCAAAT GTTCTCCAG CTGTGCCCT GAGCAAGCC AAGGCTCA

2201 TATATCTAA ACTGCTCT ACTGCTCT GAGTGTCT ACTGCTCT CTTCAGAGT CCGGAGCAG TCAATGCC ACTTACCTT GGTCTGCT  
ATATAGATT TGAAGAGC TGAAGAGC TGAAGAGC TGAAGAGC TGAAGAGC TGAAGAGC TGAAGAGC TGAAGAGC TGAAGAGC

2301 TACGCTGCT AACTTGAACA AGTAGGCTT CCGCTGAGG TTGATGAGG TGAAGGAGC ATGATGAGG CCGCTGAGG TGAAGGAGC AACTTGAACA  
ATCCAGACA TGAAGCTGT TATTCAGAA GCGGCTGTC AACTGCTC ACTTCCCTG TACGCTGCT AACTTGAACA AACTTGAACA AACTTGAACA

2401 TTGGGTTAC CAGGCTCTC TGGGCTCT TCTGCTCAT AGGAGTCT GGTGAGACA GGTGAGACA GGTGAGACA GGTGAGACA GGTGAGACA  
AAGGCTGTC GTGCGGCTG AAGGCTGTC AAGGCTGTC AAGGCTGTC AAGGCTGTC AAGGCTGTC AAGGCTGTC AAGGCTGTC AAGGCTGTC

2501 CAGGCGCAC CAGGCTCTC CATAGATTA CTTGCGGAG CAAAGGAGG CCAATCTCT AGATGCTCA AGGCTGCAA GGTCTTCA CTGACAGTT  
CTGCGGCTG GTCTGAGAG GTATCTCAT GGTCTCTC CAAAGGAGG CCAATCTCT AGATGCTCA AGGCTGCAA GGTCTTCA CTGACAGTT

2601 CTGCTCTG AACTCAGG GTGCTTGT CAGTGTCT GGTCTCTG ATCTCTCT GTTCTCTC TTGAGGCTT GGTCTCTC GGTCTCTC  
GAGGAGAGC TTGAGTCT CAGGAGCTA GTGCGGAG GTGCGGAG GTGCGGAG GTGCGGAG GTGCGGAG GTGCGGAG GTGCGGAG GTGCGGAG

2701 AGCATCTAG AAGGCTGTC CTTTGAAGT GGTCTCTC CAGGATTTA GTTCAAGG TGAAGGCTT TCTGAGGCTT CCAAGGCTT GTTGTATTT  
TCTTGAATC TTGCGGAGC CCAATCTCA GGTCTCTC CAGGATTTA GTTCAAGG TGAAGGCTT TCTGAGGCTT CCAAGGCTT GTTGTATTT

2801 TTCTGCTCT CTTGAGGCT GTTCAAGG GTTCAAGG GTTCAAGG GTTCAAGG GTTCAAGG GTTCAAGG GTTCAAGG GTTCAAGG  
AAGGAGCA GAGGCTGAT CAGTATGAG GTTCAAGG GTTCAAGG GTTCAAGG GTTCAAGG GTTCAAGG GTTCAAGG GTTCAAGG

2901 GTGAGGCTT CAGGAGGCT CAGGAGGCT CAGGAGGCT TCTGAGCTA CTTGAGGCT TCACTGCTC CATGAGGCT CAGGAGGCT TCTGAGCTA  
CAGTCCGCA CTTGAGGCT CAGGAGGCT CAGGAGGCT CAGGAGGCT CAGGAGGCT CAGGAGGCT CAGGAGGCT CAGGAGGCT CAGGAGGCT

FIG. 3C-1

3001 AGTCAAAAGC TGTTCACCT CACACTTCTT OCTGCCATG AAAGCTGAG TGCAGCGGC CTTCATCAC TGTATGGAG CGAATATCAG GTTTCAGCC  
TCACTTTTC ACGATGTGA CTCTAAGAA GCGCCGTC TTTGGACTC AGCTCGGCG GAGGTACTG ACATAGCCTC CCTATATGC CAATGTCCG

3101 CAATCEAGCG CAGCTGCCA AGCTGACTT OCTGAGTAC CCAACATGC OCTGCCAGC CTTCGTACG CAGAGAAAC CATGCCAGC GCGCTAGAT  
GTAGATGCC GTGACGGGT TGCAGTGA GCGATCCAT GGTGTATAG CGAGGTTGT GAGCCAGTG GTCTCTTTG GTACGCTGT CCGATCATA

3201 GAAAAAGGC CTCAGGGTG CCATGCCAG OCTCTAGCC AGGCGTTTG CAGCTGGGC GCGAGCTTC TCGAATCTG CTGTCTGCC TGAATAAGA  
CTTTTTCCT GATGCCGC GTAGCGTCC GCGATCGCG TCGCGAGGC GTTGGAGCG GCGCTGGAG ACCTTAGAG CTGTCTGCC TGAATAAGA

3301 AGCAGACTG AGAAGATTC CTAGTTCCT GGTTCCTGC CTTTATTC CTATCTCTT GCGCCAGCC CATTCGCC TCCTAACAC AGCTGAGC  
TGTCTGACT TCTCTCAG GATCAGGCA CCGAAGAG CCGAATTAAC GATCAGGCA CCGGTCGCG GATTCGCC TCCTAACAC AGCTGAGC

3401 AAGGTCACA TTCCAGAAC CCGCCGCCA GCGAGCTTG GAAACGAA AGCTGCCCA AGCCAAAGT CAGTAGGTC AGCGCAGC GCGATACAC  
TTCCAGTGT AAGGCTCTT GGTGCCGT CTCTCGACC CTTTCTCTT TCGAGCGGT TCTGTCTCA GTATGCCAG TCGCGCTCT CCGATATGC

3501 GCTAGCTTA GCTGGGAGG TCGAAGAG CAGTGTGT CAGGCTCTG CCGATGCCG TGAATCTGC TGAAGCTAC TTTTATTA GTGGAGCAT  
CGATCGAAT GCGGCTCTC AGCTTCTTC GCGACGACA GTGGAGACT CCGTCCAGC AATGAGCG ACTCGGATG AAAATATTT CAGGCTCTG

3601 GGTGCTTGC CTATCAGT GTTTCAGAT TCGGTGAGT AGAAGAGCA AAAGCTTTG TCGGTGAGT AGCTTCCAC TCATTCCTT AAGGCTAT  
CGAGGAGC GATGAGTCA CACTCTCTA AGGCTCTGA TCTTCTCTT TTTGAAAG AGGAGCTCA TCGAGGTTG AGTAAAGTA TTGCCAATA

3701 GATTTACTG TTTATCAG CTAGTCTCT GTCCATCTT AGGCGGCT TCGAATCTG ATTTTGGCG CAGAGAGCG GGTGGGCG GAGCTGCCA  
GTAATGAC AAGTATGTC GATCAGAA CAGGTAAGA TGGGGGCG AGCTTAGAC TAAAAAGCG GTTCTTCCG CCAAGGCTCT CTGAGCGTT

3801 GCACTTTGG GAGGTTTTC TTTCTCTC AAGAGAAC AAAGCTTCT TCTGGGCTC TCTTCTCT CTCTAGCTG GGTGTACAG CAGTAGAGT  
GTGAAAGC CCGCAAGG AAAGAGAG TATTTCTTC TTTGAGTA AAGAGCGAG AGGAGAGA GAGATTGAC CCAAGTCTC GATCTCTCA

3901 AGTGGCTAG AGCTATCTT TCTTCTTA TTTTCTTG ATTTATAT TTAATTTT GTTATAGT GTCTCTCAC ATGTGATCT GTGAGCACA  
TCAAGATG TCGATAGA AGAAGAAAT AAAAAATC TAAATATA AATACAAA CACATATCA CAGAGAGTG TACAGTAGA CAGGTGTGT

4001 TCAATCTTT GTTCTATG AGCTCAGAG AGGCTTTCA ATAGCTGCA ACTGAGTTT TGAAGTTA TCACTGCGG TGTGATCT CAGATCTAA  
AGTACAGA CACATATC TCGATCTTC TCGGAAAT TATGAGCT TCACTCAA ACTGTCAAT ACTGAGCGC ACAGCTAGA CTCTAGTTT

4101 CCGAGTCTT CTGAGAAC AAGTCTCTT AAAGCTGAG CACTCTTTC AGTGCAGAG CCGATCTCT AGGCTTTC TATGATTT ATCTGGGG  
GGTTCAGCA GACTCTTTG TTAGAGAA TTTGGACTC GTAGAGAG TCAAGGCTC GGTAAAGAC TCGAAGTG AATAGTTAC TATGAGGCG

4201 GAGCAGCTG GCGACCTTT CATGACTTC ATTTATTTT AAAAAAAT GCACTCATG GCGATCTTT CTAGACTAC ATCTAAGTG GATTTCTCT  
CTGTGGAC CCGTGTGAA GTTCTGAG TAAATAAAT TTTTCTTT CCTGATAC CCGATGAA GATCTGAGT TATGATTCAC CTAAAGAGA

4301 AAGAGAGT CTCTAGCG GTAGAGTGC AGTTTTCG CCAATTTCA AGCTGCGA CACTCTGAA GCGGCTCGT TTTCTTTCT GTATTCAG  
TATTTCTCA CAGTGAAG CATCTAGG TCGAAGGC GGTTCAGGT TGTGAGCTT CCGGAGCGA AAGAGAGA CATGATCTC

4401 GCGAGCTGC CTTCGTCTT TCTCTCTAT GCGGCGAGT AGTCTAGAG CCAATGAA ACCTAAAT TACTTCTTA CAGAGCGTG AAGCTAAGT  
CGTGCAGC GAAAGCAG AGAGAGATA CTTGGCTCA TCGAGTGC CTTTCTCT TGTGATTTA AATGAGGAT GTCTGCGAC TTGATTTCA

4501 GCAAGCGGC ATTAAGCG TTAGAATC TCACTGGA TCTTTAAC ATCGGAGCG GAGGTGATA CATGAGCA GCTTCTTTC ACATTTTGG  
CCTTGGCG TATTTTCG AAATCTAG AGTTGAGCT AAGAAATG TAGGCTGCC CTGACCTAT GTACATGCT CCGAGGAGG TGTAAAGCC

4601 GAGCGAGCG AGCGTAGCA AATGAGAC AGCTCTTTC AGGCTTTCT ACAGATCTT GCGACGAGC AAGGAGAG TCGGAGAG AGGCGAGCG  
CTGGCTGC TCGCATCT TCACTCTG TCGAGATG TCGGAGAA TGTCTGAA CCGTGTGT TTTGCTCT AGGCTCTCT TCGGCTGCG

4701 AGTGTGCG GTGGTGGAG AGCTGAGTA GCTTGGCG TCGTGGCG GCGAGCGCG TGAAGCTAG AGGCGCGCG TCAATCTTT GACTCTCTG  
TGCAGCGG CAGGAGCT TCGAGCAT CCGAGCGCG AGGCGCGCG CCGTGGCG ACTTTGATC TCGGCGCG AGTTAGAA CTGAGAGAC

4801 CTCAGAGCG TGTGTCTGT TGAAGCTT AGCTGCGTG TCTGAGAT GAGAGCGCG TTTTTCGCG CCGGCGGT CTCTAGCTC CCGGTCTGT  
GATCTGCG ACGAGGCA ACTGAGAA TCGAGCGAC AGAATCTA CCGTCTCG AAACAGCG CCGGCGCG GAGATGAG GCGGAGAGC

4901 TCACTCTT TCTCTCTT CAGGCTTC CTAGTCTCT GATGCGCG CCGGCTGC TCTCTCTT TCTACTTT TAGGCGCA CTTTACCG  
AGTACGAG AGAGAGCA GCGGCGAG GATTCAGCA CTCAGGCT CCGGCGAG AGGAGAGA AGATGTGAC ATCGGCTGT GGAATGCG

FIG. 3C-2




FIG. 4A



1 GCAGCTGACC AACCTTTGCG GATGCGGCTG CAAAGTATAT ACCGCGTGGT TACGAGAGCG TCAGAACTTT TACCGGAAAG CGGCTGCTCT AACCGGAGCG  
 CTTGAGCGCG TTTCGAGCG CTAAGCGGCG GTTTCATATA TGGGCGACCA ATGCTCTTGG ACTCTTGAAA ATGCGCTTTC GCGCGAGCGA TTGCGCTTGG

101 TACCGAAGTA GCGGAGAAA AACGAAAAA AAATTCGAGA GAAGCTTCCA GAGCTGCTG CTCTTGGCTG TTCTTCAAA AGGACTGCAA GTGCGGAGTC  
 ATGCGTTGAT GCGCTCTCTT TTCTTTTCTT TTGAGGCTCT CTTGAGAGGT CTGCGAGCGG GAGGAGCGAG AAGGAGTTT TCGTGAAGTT CAGCGGCTCG

201 ACCTGCGACG CAGCAGAGGT TACGCGCTG AACCGCGCTG ACCTGCGCTG GCGCTGCTG CCAAGCTAAC GCGCGACCGG TCGGTAAAGC GTGACCGGCT  
 TGGGAGGTGG GTGCTTCTCA ATCGCGAGCG TTGCGCGCGG TCGCAGCGCG GCGCGAGCGG CTTCGATTG CCGCGTGGCG AGCGATTG CACTGCGGCA

301 CCAATCTCTG GTCTGAGCGG GCGCAGCGCG AACCGCGCGG ACCTGCGCGG GCGCGCGCGG TCAAGTGGCG GCACTTTCTA TTTCGAGAGC CCGAGCGCGC  
 CTTTACGAGG CAGCTGCGG CCGCTGCGGT TCGCGCGTGG GCGCGCGCGG ACTGCGAGCG CCGTGAAGAT AAGGCTTTC GCGTGGCGTG

401 GTTCTGAGCG GAGCGCGCGG CCGCTGCTTG TGATTGCGTG TCGCGCGCGG CTTCGAGCAA TCAAGCTTCC CTTCGCTATT GTGAGCGGTA CCGCTGCTG  
 CAGCGATTG CCGCGCGCTG CCGCGGAAAC ACTGAGCGCG AACCGCGCTG GAATCGGTT ATGCGCGAGG GAGGAGTAA CATCTCGAT CAGCGGAGCG

501 TTCTTTTCTG TCGTCTTCTG CCGCTGCGTG GTCTGAGAGA GCGAGCGCGG GATCTTTCTG GCGATGGCGA CTGCTTCTCA CCGCGGCTG TCGGAGGACT  
 AACGAAAGC AACGAGAGG CCGCGCGCGG CAGAGCTTCT CCGTCTGATC CCGAGAGAGG CCGTGAAGCT GAGCGAGAGT GCGGTGAGCG AACCTGCTGA

601 TGTGCGAGCG TCGTCTTCTG TCAAGAGCTA GAGCTTTCTG CCGTGAAGTT AACCGCTCTA AACCGGAGCT GAGCGGCTCA TCGTCTTCTG CCGAGAGCTA  
 ACAGAGCTGG AACGAGAGC AGTATTCGAT CCGCGAAAGC CCGTCTGAGT TCGCGAGCTA TCGCGGAGT CTTGCGAGT AGGAGAGCT CCGTCTGCTA

701 AGTCTCTCTG GAGCTCTGAG CACTGAGTCC TTGCGAGGTT TTCAAGAGCT AGTGAAGCTA GCGTGAAGCG AACCGCGCGG GTGCTCTCTA AAGGCTCTG  
 TCGAGAGGTA CCGGAGATCT GTGAGTCAAG AACCTGCGAG AAGTCTTCTA TCGTCTCTG TCGGTGCGTG CCGCGAGAT TTTCGAGAGC

801 CTTGCGTGGG TTGCGAGGCT CTGATTGCGG AACGATTCTG CCGCTGCTG CCGCGCGCGG CCGTGGCGG GATCTCTCA ATGCGAGAGT CCGTGAAGT  
 GAGCGGATC AAGGTTGCGA CACTGAGCGG TCGTGAAGT CCGGAGCGG CCGTGGCGG GATCTCTCA ATGCGAGAGT CCGTGAAGT AAGCGCTTCC

901 GCGCGCGCGG GCGCGGCTG CAGCTTTCTT GCGCGCGCGG ATGCTATCTG AACCGATGCG CTGCGAGAGA GTCTGAGAGA TTCTGCTCT CACTTTTCTG  
 CCGCGCGCGG CCGCGGCTG CCGCGAGAGT CCGCTGCGTG TCGAGAGAGG TCGGTGAGG GAGCTTCTG CCGAGCTCTG AAGGAGAGT GTGAAAGAG

1001 TCGCGTGGG TTCACTGAAA CCGTCTGCTG TCACTGCGAA GAGAGAGAGA CTGCGAGCGG ATGCTCTCTA CTCTGAGCGG GTTGGCGGAA CAGCGTTGCA  
 ACGAGAGTCT AAGTGAAGTT GCGAGTCTG AGTGAAGCTT CTCTGCTCTG GAGCGTTGCG TCGAGAGTT GAGGATCGG CCGCGGCTG GTGCGAGCT

1101 ACTGCGAGCT GCGCGCGCGG GAGCGCTCTA CCGTGGCGG GTGCTTTCTG GTGCGAGTGG AGGAGGAGCG TTGCGGAGG CCGTCTGCTG TCGTCTCTG  
 TCGCGTGA GCGCGAGCGG CCGCGAGTCT CCGTCTCTG CAGCTCTG CAGCTCTG CCGTCTCTG AAGCGATTG CCGCGAGTCT AAGCGAGTCT

1201 TCGTCTGGG GTTGGCGGCT TTGCGCTCTA TCGTCTGCTG AATGCTCTG CCGCGAGTGG CCGTCTCTG CCGCGAGTCT TCGTCTCTG CCGTCTCTG  
 ACGCGAGCGG CAGCGCGCGG AAGCGAGAT ACGAGAGAGC TCGAGAGAGC CCGTCTCTG CCGTCTCTG CCGTCTCTG CCGTCTCTG CCGTCTCTG

1301 CCGCGCGCGG ACGCGGAGG GTGAGAGAG CCGCGTCTG TTGAGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 CCGTCTCTG TCGAGAGAG CAGTCTCTG CCGCGAGAG CCGCGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG CCGTCTCTG CCGTCTCTG

1401 GTTGAAGAGA ATCTCTCTT AACCGAGTGG TCGAGAGTGA GTGAGAGAGT CCGTCTCTG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG  
 CAGCTCTCTT TCGAGAGTGA AACCGAGTGG TCGAGAGTGA AACCGAGTGG TCGAGAGTGA AACCGAGTGG TCGAGAGTGA AACCGAGTGG

1501 TCGTCTCTG CAGAGAGTGG GTCTCTCTA TCGTCTCTG CCGTCTCTG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 ACGAGAGTGA GTGCGAGAG CAGAGAGTGA AACCGAGTGG TCGAGAGTGA AACCGAGTGG TCGAGAGTGA AACCGAGTGG

1601 GTGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

1701 CTGAGTCTT CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 GAGTCTCTG CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

1801 CCGTCTCTG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 CCGTCTCTG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

1901 CCGTCTCTG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 CCGTCTCTG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

2001 TCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 ACGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

2101 GAGCGAGTGG TCGAGAGTGG CCGAGAGTGG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 CCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

2201 AAGTCTCTG AATTTCTCT CCGAGAGTGG CCGAGAGTGG CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 TCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

2301 TCGAGAGTGG AAGTCTCTG CCGAGAGTGG CCGAGAGTGG CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 AAGTCTCTG AAGTCTCTG CCGAGAGTGG CCGAGAGTGG CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGTCTCTG

2401 AAGTCTCTG AAGTCTCTG CCGAGAGTGG CCGAGAGTGG CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 TCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

2501 CCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 GAGTCTCTG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

2601 CCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 GAGTCTCTG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

2701 ATGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 TCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

2801 CCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 GAGTCTCTG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

2901 CCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 GAGTCTCTG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

3001 GAGTCTCTG AAGTCTCTG CCGAGAGTGG CCGAGAGTGG CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 TCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

3101 CCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 GAGTCTCTG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

3201 TTCTCTCTG AAGTCTCTG CCGAGAGTGG CCGAGAGTGG CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 TCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

3301 TCGAGAGTGG TTCTCTCTG CCGAGAGTGG CCGAGAGTGG CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 AAGTCTCTG AAGTCTCTG CCGAGAGTGG CCGAGAGTGG CCGTCTCTG CCGCGAGAG CCGTCTCTG CCGTCTCTG

3401 CCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 GAGTCTCTG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

3501 CCGAGAGTGG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG CCGTCTCTG  
 GAGTCTCTG CCGAGAGTGG CCGCGAGAG CCGTCTCTG CCGCGAGAG CCGTCTCTG TCGAGAGAG CCGTCTCTG CCGTCTCTG

FIG. 4B-1







10801 GTCTGAAA AAAAAGCA GAGAGGAG TCGAGGCA ABACTBA CATTCTGT GTGTCTTT CTGAGTCA TCTGAGAG CAATCTGC  
CAGACTTT TTTTCTCT CACTGCTT ACTCTGCT TATGAAAT GTAGAGAC GACAGAAC GACATCAT AACTATTC GTAGAGCC  
10901 TTCTGCAA GTGAGAGT AACATTCT TBAAGAGT ATTGCTCT CTGATTTT GTTTTGT TATGCTCT AGATGAGC CAGAGGCT  
AAGAGGTT CATCTTCA TTGAGACA AATTTTCA TAAAGAGC GAAAGAAA GACAAATA ATGACAGC TCTAGCTT GTCTGAGA  
11001 TCGAGCAA GCTAGCTT TTAGCTCA GCGAGCTC AGCTTCTC TGGGATTC TACGAGAG TTCTAGCT GAGGAGCT CCGAGGCT  
AGGTTGTT CCGTGCAC ATGCTGCT CCGTGTAG TCGAGGTT AGGCTGAG ATGCTTCT AAGATGTA CTTGTTCA CCGTGGCT  
11101 ATGCTCTT CAGATTTT AGGAGTCT ATGCTGCT TTGATCTT TACAGGCT TACGAGCT TCGTT  
TAGGAGAG CTTCTAGA TCGTCAAG TATGATGG AACTAGAA ATTCTGAG AATGCTCT AGTCA

FIG. 4B-4




FIG. 5A




FIG. 5B



<b>VEGF project</b>	<b>VEGFR2 project</b>	<b>Tie2 project</b>
<b>Screening primers</b>	<b>Screening primers</b>	<b>Screening primers</b>
Primers: VF1-VR1A Product size: 0.69Kb	Primers: KF1-KR1 Product size: 0.45Kb	Primers: TF3-TR1 Product size: 0.45Kb
<b>PCR program</b>	<b>PCR program</b>	<b>PCR program</b>
Hot start	Hot start	Hot start
94°C 40 sec 65°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 58°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 58°C 1 min 30 sec 72°C 1 min 30 sec
40 cycles	40 cycles	40 cycles
<b>Confirmation primers</b>	<b>Confirmation primers</b>	<b>Confirmation primers</b>
Primers: VF2-VR2 Product size: 0.98Kb	Primers: KP2-KR2 Product size: 0.58Kb	Primers: TF2-TR1 Product size: 0.47Kb
<b>PCR program</b>	<b>PCR program</b>	<b>PCR program</b>
Hot start	Hot start	Hot start
94°C 40 sec 65°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 65°C 1 min 30 sec 72°C 1 min 30 sec	94°C 40 sec 58°C 1 min 30 sec 72°C 1 min 30 sec
40 cycles	40 cycles	40 cycles

*FIG. 6*





FIG.7



*FIG.8*





FIG.9



**FIG.10**

## SEQUENCE LISTING

<110> XENOGEN CORPORATION

<120> TARGETING CONSTRUCTS AND TRANSGENIC ANIMALS PRODUCED  
THEREWITH

<130> PXE-008.PC

<140>

<141> 1999-12-16

<150> 60/152,522

<151> 1999-09-03

<160> 39

<170> PatentIn Ver. 2.0

<210> 1

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer PGKF

<400> 1

atcgaattct accgggtagg ggaggcgctt t

31

<210> 2

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer PGKR

<400> 2

ggctgcaggt cgaaaggccc ggagatgagg

30

<210> 3

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer NeoF

<400> 3

acctgcagcc aatatgggat cggccattga ac

32

<210> 4

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer NeoR

<400> 4

ggatccgcgg ccgccccag ctggttcttt ccgcctc

37



<210> 5  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer TKF

<400> 5  
ggatcctcta gagtcgagca gtgtgggtttt 30

<210> 6  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer TKR

<400> 6  
gagtcctcgt agtcagggtt agttcggtccg 30

<210> 7  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer F5R51

<400> 7  
gtacatttaa atcctgcagg 20

<210> 8  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer F5R52

<400> 8  
agtcctgca ggatttaa 20

<210> 9  
<211> 77  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer F3R31

<400> 9  
ggcccgggct taattaatgc atcatatggt accgtttaaa cgcggccgca agcttggtcga 60  
cggcgcgccg gccggcc 77

<210> 10  
<211> 77  
<212> DNA  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer F3R32

&lt;400&gt; 10

gatcggccgg ccggcgcgcc gtcgacaagc ttgcggccgc gtttaaaccg taccatatga 60  
tgcattaatt aagcccg 77

&lt;210&gt; 11

&lt;211&gt; 39

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer VN1R

&lt;400&gt; 11

ctgtatttaa atctgcccac cctattcagg acagtagtc 39

&lt;210&gt; 12

&lt;211&gt; 31

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer VN1F

&lt;400&gt; 12

ccaatgcac aaccagcca ggaggagtgc g 31

&lt;210&gt; 13

&lt;211&gt; 38

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer VN2R

&lt;400&gt; 13

aacgcgtcga cttcgagat gtttcgggga taaccagg 38

&lt;210&gt; 14

&lt;211&gt; 40

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer VN2F

&lt;400&gt; 14

ttggcgcgcc ccatagagaa gagacaccaa aggcacgctc 40

&lt;210&gt; 15

&lt;211&gt; 36

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer FosB1F

&lt;400&gt; 15

ctgtatttaa atcccgtttc tcactgtgcc tgtgtc 36

<210> 16  
<211> 35  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer FosB1R

<400> 16  
gtctcctgca ggcttcctcc tccttggtcc ttgcg 35

<210> 17  
<211> 35  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer FosB2F

<400> 17  
aacgcgtcga cggatgggat tgacccccag ccctc 35

<210> 18  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer FosB2R

<400> 18  
ttggcgcgcc ccttgctcc acctctcaaa tgc 33

<210> 19  
<211> 31  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer VF1

<400> 19  
acctcactct cctgtctccc ctgattccca a 31

<210> 20  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer VR1A

<400> 20  
gctctggcgg tcacccccaa aagca 25

<210> 21  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer VF2

<400> 21  
ccctttccaa gaccgtgcc atttgagc 28

<210> 22  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer VR2

<400> 22  
actttgcccc tgccctctc tctgttcgc 29

<210> 23  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer KF1

<400> 23  
gctgcgtcca gatttgctct cagatgcg 28

<210> 24  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer KR1

<400> 24  
ttctcaggca cagactcctt ctccgtccct 30

<210> 25  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer KF2

<400> 25  
cagatggacg agaaaacagt agaggcgttg gc 32

<210> 26  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer KR2

<400> 26  
gaggactcag ggcagaaaga gagcg 25

<210> 27  
<211> 29  
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer TF3

<400> 27

agcttagcct gcaagggtgg tcctcatcg

29

<210> 28

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer TF2

<400> 28

caaatgcacc ccagagaaca gcttagcctg c

31

<210> 29

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer TR1

<400> 29

gctttcaaca actcacaact ttgcgacttc ccg

33

<210> 30

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F51

<400> 30

cccagtgtct ctgatttagg gagagcacct gag

33

<210> 31

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer R51

<400> 31

ccagactgcc ttgggaaaag cgcctc

26

<210> 32

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer F52

<400> 32

cagtgagagt cttctctgtc cctcaatcgg ttctg

35

<210> 33  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer R52

<400> 33  
tggatgtgga atgtgtgcga ggccag 26

<210> 34  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer F31

<400> 34  
aatcaaagag gcgaactgtg tgtgagaggt cc 32

<210> 35  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer R31

<400> 35  
cggctcccca aaatgtggaa gcaagc 26

<210> 36  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer F32

<400> 36  
gaatccatct tgctccaaca ccccaacatc 30

<210> 37  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer R32

<400> 37  
cgcctectct cccagtcctc cccttg 26

<210> 38  
<211> 4998  
<212> DNA  
<213> Mus sp.

<400> 38



tccacccacc	tgtttctcac	gtccccggcc	ttcctagtta	acttcatggt	taaagaagcc	60
tcacccgggg	agggtgtggt	gccacagaag	gaagggtgct	cccacaagcc	cccagtgtct	120
ctgatttagg	gagagcacct	gagccagtg	agagtcttct	ctgtccctca	atcggttctg	180
aaattcccca	cttgccctcc	ttatccaggg	gacagggtg	cccaccctat	tcaggacagt	240
agtcttaaac	tcgtagccaa	cagacttttt	attgggctgg	gagaaagaga	tgaggctcct	300
gaagctcage	cgagtgggct	ctgattccta	cttctcagag	gtcgggcagc	ccagccaata	360
ctgagcaatg	gagcgtgggt	agggaggatt	cacagagtcc	actcgccggg	ttctaagggt	420
gactcggtag	tatttgtctg	aaagaaagaa	tggaaaaagg	gttatgtgag	attctgcctg	480
atcctgtcca	ctgggtcccaa	gaaggataaa	ggctttttct	cagaaggga	agtgaacatc	540
caccaagcag	ataatgtcac	catctacagg	ctgtgttcag	caccagggga	ccaagacctg	600
caggcaaggc	ctagccaaaa	ccagtctaag	gagtagaaag	gggctcccac	ctccagagaa	660
gaaatagacg	ctctgaatgg	gctcgcaggt	ggcaggtaca	agccagtcca	tatcataatc	720
atagttgttg	taggttccta	gccactctc	ctcgtggag	aacaaagaga	accagattga	780
acgtgatgaa	cgacgggagt	tcgagctctg	gctgcgtctg	tggccacgcc	ctcggcgtga	840
acgatagcgc	tttcggcttc	tacgcttaga	cttctgtttt	ttggcttggg	cagagtggga	900
taaggagcca	gtgacgtaga	tgcggccggc	catagcagcg	tccactttcc	ctggcacacc	960
atgccagttc	cggctgatga	attgggggtc	tctggctcca	tctgtaacag	ggaaggggtt	1020
aatgcacttg	gcagattctg	gctttgattt	ctccagcaag	gttgtctgtc	tatctattta	1080
tctatcttta	tctatgtatc	tatctatata	tctatgtatc	tatctatcta	tcctctacct	1140
acctacttac	ctatctatgt	atctatctat	ctatcatcta	cctacctact	tacctatcta	1200
cctattttatt	tgtttgtttg	ttttctttga	aacaggatct	tagcacctac	ctatggctgg	1260
tttgcaactc	actatgaagc	cataactggc	ctcttaactc	acaaagatcc	acttgccctgt	1320
gtctctgagt	gctgggatta	aaagcatgtg	ccactacacc	cagctccagt	aggaccttta	1380
gaacacattt	gctatgcctt	gcctaagaca	cacaactcag	tccccaggcc	ccagcctccc	1440
tgtctagagc	tttttcccat	cctctctcca	ctgtatccct	tgaatctctg	ccccatccga	1500
aaccctcag	cgcgcagccc	ctccttctgc	tgtgttaggc	aaagtccaag	gtatgggata	1560
caaatagagc	caagcctcat	ccccaaaag	tcaacagaag	caaagtctag	ccagagcaaa	1620
cagctcttga	tcgatggtgt	cacagttcca	ggccctctcc	ctggaagccc	ccactatcac	1680
agcccagttt	ccagagaaaag	aagccagcct	tgctctccct	ccataccaga	ggatctgccc	1740
cagaagagga	gttcgaaaat	gttctcccag	ctgtcccgt	gaagcaaggc	aaagtgtcta	1800
aacacggctg	acagagagct	gccttcgcac	tcctcctggc	tgggttgetg	ctgaaattcg	1860
tactcccagt	actgcttccc	tgaggagcag	aacagctggc	atcaggagag	atctgaccaa	1920
ggcagagagg	aatcatggaa	tagaacaggg	actccaccac	ctgccccctt	ctcctccacc	1980
ctgagtaccc	ttgaagaagt	agacccttcc	ccggccactg	taacgggtggg	caggaagggc	2040
gaacgctgca	tcaacattgt	ctggtatgcc	actgaagcct	tcggagatgt	ttcggggata	2100
accagggtcc	aggaccccat	cctcaaagcg	ccagtactga	ctaccctgaa	agacagagat	2160
cagaaggggtg	aggacatacc	gctggccaca	gaagcagtc	tatatcctaa	actggctgtc	2220
acctgctcct	ggagtccctg	actgctttgt	cttcacagct	ccccagcacg	tccatggcac	2280
cctttacctt	gcctcagact	taggtctggt	accttgaaca	agtaggtctt	cccctgacag	2340
ttgatgcgag	tgaaggcagc	atcgatgggg	ccctcaatgc	cccagacatc	ttggataagt	2400
ttgggggtacc	caggcctcac	tgccgtctca	tctagctcat	agcagtactg	ccctagaaca	2460
ggggaaactg	tgtgagaagc	agatgagcct	aaggcagatc	cgaccgccac	cagacctgtc	2520
catagagtca	cctcggaagg	caaagaggga	cccatctctg	agatccgtga	aggcgtcaaa	2580
gggctttcca	ctgcacagtt	cttctctctg	aaactcaggg	gtcccttgat	cagtgggtgtc	2640
gggccttagg	atctcctcct	gttgctccac	tttaggcgt	ggggtgcttg	gctgttcctc	2700
aggatctagg	aaggctgtcg	gctttagagt	gccgtccgtc	cgaggattta	ggtcaccggg	2760
tggagagggtg	ttctcgggtt	gcacaccggg	gttggtattg	ttcttgggct	cctccacgta	2820
gtcatagctc	caataatcat	cctctggcat	agtgaacacg	tccccccgcg	ttactgcagg	2880
cagaacgggg	agcagtgagt	gtcaggctgt	ggaggggagc	ccaggcccac	ccaccagggc	2940
tctgaactca	ccttggggct	tgcactgtct	catgtagtctg	gcacagcagc	tctgatagta	3000
agtgcaaagc	tcgtcacact	gacacttctt	gctggccatg	aaaccctgag	tgcagcggcc	3060
cttgcatgac	tctatgggag	ggaatatcag	gtttacagcc	caatctaggg	cacctgcca	3120
acctgcactt	ccctaggtac	ccaccaatcc	cctcccacac	cttggtcagc	cagagaaacc	3180
catgccacca	gggctagtat	gaaaaagggc	ctcagggggtg	ccatggcagg	cctctagccc	3240
agggccttgg	caagctgggc	gcggagcttc	tggaaatctcg	ctgtcctgcc	tgaaaaaaga	3300
agcagactga	agaagagttc	ctagttccct	gggtttctgc	cctttatttg	ctcatcctct	3360
ggcccagccc	cattgccctc	ctccaaacac	agctgcagca	aagggtcaca	ttcccagaac	3420
cccagcccca	ggagagctgg	gaaacagaaa	accctcgcca	agaccaaagt	cagtagggtc	3480
acgggcagga	gggataaac	gcttagctta	gctggggagg	tggaaagaag	catgtgttgt	3540
cacctctga	gccagtcctg	ttaatctccc	tgagccttac	tttttataaa	gtgggacccat	3600
gggtgccttg	ctcatcaggt	gttgagagat	tccgtgagct	agaacagaca	aaacgtttcg	3660
tgctggagc	agcttccaac	tcattcccat	aagccgttat	cgatttactg	tttgatcagg	3720

ctagggtgctt gteccatcct acccccgcct tcgaatctgg attttttgggg caagaagggg 3780  
ggttggggga gagctggcaa gcactttggg ggaggttttc ttttcttctc ataaaagaac 3840  
aaagcttcat ttctggcctc tccttggtct ctctaagctg ggtgttacag cataggaagt 3900  
agtgggtcag agtctattct tctttcttta ttttttttag atttatttat tttatgtttt 3960  
gtgtataagt gtctgctcac atgtgcatct gtgcaccaca tgcagtctct gtgtctatgg 4020  
aggtcagaag agggctttga ataccctgga actggagttt tgaacagtta tgagctgccg 4080  
tgtggatgct gagaatcaaa cccaggtcct ctgtaagaac aagtactctt aaaggctgag 4140  
ccatcttttc agtcccagag cccattcctg aggctttcac taatccattg atcctcgggg 4200  
gaccaccctg gccacacctt caatgacctc atttatttta aaaaaaaat ggactcattg 4260  
ggcatacttt ctagactcac atactaagtg ggatttctct ataaagaagt gctcactggg 4320  
gtagagtgcc aggttttggg ccaaattcca agcactggca cacttctgaa gcccctcctg 4380  
tttctgttct gtaatcacag gcgagcgtgc ctttggtgtc tcttctctat ggaccgcagt 4440  
agtctcagcg gcaaaatgaa acactaaatt ttactccta cagacgcgtg aagcctaagt 4500  
ggaaaccggc attaaagggc tttaagaatc tcaactgcga ttctttaacc atccggaggg 4560  
gacgtggata catgtagcca gcttgcttcc acattttggg gagccgagcg agcggtagga 4620  
aatggaagac agctctttac agcctttctt acagcatctt gcacaccacc aaggggagac 4680  
tggggagagg aggcggagcc aggtgtgggc gtggctggag acctggggta ggcttgcgcc 4740  
tgcgtcgggg gcggagcccg tgaaacctag aggcggggcg tcaaatcctt gactctgctg 4800  
ctcagaggcg tggttgctgt tgagcatctt agctccgctg tgcttagatt ggagcagcgc 4860  
tttgttccgg gcaccggcgt ctctaccctc ccgctctggg tccatgcttc tctctccctt 4920  
catgcccttc ctaagtcgct gagtcccgga gctgccctcc tccttctgct tctacacttg 4980  
tagccagca cctttacc 4998

&lt;210&gt; 39

&lt;211&gt; 11176

&lt;212&gt; DNA

&lt;213&gt; Mus sp.

&lt;400&gt; 39

gcagctgggc aaacgttggc gatgccgggtg caaagtatat acccgggtggt tagcagaagc 60  
tgagaacttt tagccgaaag ccggtccct aagccgaagc taggcaagta ggggaagaaa 120  
aagaaaaaaa aaattccaga gaagcttcca gagcctctc ctcttccctc ttcttcaaa 180  
aggactgcaa gtccgcagtc accctccacc cagcaagagt tagggcctcg aaccctgggtc 240  
acgctgcctc cgctcctgc cgaacgtaac gggggaccgg tgcgtaaagc gtgacgcgct 300  
ggaatcctcc gtctgacgcg gggcacgcac aggcgcagc ccctccgccc gcccgcgcc 360  
tgacgtccgg gcacgttcta ttttggaacg ccgaggccac gttgctaagg gagggggcag 420  
cgtggctttg tgattggctg tcgcgcgcag ctttagccaa tcagcgttcc ctctctattt 480  
gtagagcgta gctcccttcc ttgctttttg tggttcttcc cgtgctgggg gtctccaaga 540  
ggagagctag gattcttgtc gcgatcggga ctcgttgtca cccatgggtc tgcgaggact 600  
tgtgtggacc tggctctgtg tcataagcta gaggttttg gctgagtgtt agcgcctcta 660  
agggggaact gaaggcctca tccttctcag gcacacatat acgtgctcct gagctctaga 720  
cactcagtc ttccgaggtg ttcaaacact agatgagcta gcctacggag aggcagccag 780  
gtggctctta aaaggtctgc ctcccttag ttccaggt ctgattggcc agggattcag 840  
cccttccctc gccacgcccc ctagagtagt taagcctcta ggattccact tgcgggaagg 900  
gggggggggg gggcgtgatg gacgttctt ggggacgcag atcctatgtc acccatccc 960  
ctgcaagaca gtctgagaga ttctcgctgt cacttttctc tgccatcag ttactgaaa 1020  
cctgtcagtc tcaactggaa gagacagaca ctcggaaggg atgctctcaa ctcttaggcc 1080  
ggtcccccaa caccgttggg actgggatct ccgctgcgg gagcctcat gcagtggggg 1140  
gtgtgtttgt gtgtgagtgg agaggaaggc ttggctaagg cctctccctc tccctccctc 1200  
tgtgttgggg gtgtgggggt tttggctgta tgtgtgtgtg aatgtctgtg gctccatccc 1260  
gggagtttgt caccaggttc tgtccagcct cctctccac ccaccccc acacctaaga 1320  
gtcaccaacc cggggtgtga ttaccaccc gctggaaccg tgcaacctt ccccgaggaa 1380  
gaaggaggag gtagaaggca gttgaacaga atctctcatt aaccactgcg tcacggtgta 1440  
gtggaagggt ggggtgtgtg gctttttgcc tgtgacacac acatccacac ccgctcacc 1500  
tgtgtcact cacagggtcg gtctctctta tctctcttg gcgtgtgtgt gtcgggtggct 1560  
ttgtttgtgt gtctacgcct gtgtgtgtat gtctcaccgc gtaggagtgc gccggtctcg 1620  
gggaaatgcc cggtccttc gtgccaacgg tcaccgcaat cacaaccagc caggatcttc 1680  
agtggctcgt gcaaccacc ctcatctct ccattggcca gtcccagggg cagccactgg 1740  
cctcccagcc tccagctgtt gacccttatg acatgccagg aaccagctac tcaaccccag 1800  
gcctgagtgc ctacagcact ggcggggcaa gcggaagtgg tgggccttca accagcacia 1860  
ccaccagtgg acctgtgtct gccgctccag ccagagccag gcctagaaga ccccgagaag 1920  
agacagtaag tatgaggcct caggagtgg gatggaggag cctagctagg gatgtgggct 1980

cagtttgtac	agtgccttgc	tgccatgcat	gaagatccct	agcacagcat	aagccaggag	2040
tggttatgca	gacctgtaac	cccagctctc	agaaggtgga	ggcaggagga	gcaggagttc	2100
gaggccagcc	tgtgctactt	atggagtcca	gcctgcactg	caagagatca	ttattttcaa	2160
aagttggcct	tggggggagg	tgggtgaggg	aagtaagaga	aagtgacagt	aatttttgtca	2220
cttaatagtt	ggaggttcct	ctgaggcctc	aagtctgaag	gaactttacc	attctggcca	2280
gtgaggagta	gggggttatta	tttgggggttc	aggaggaagg	aagttttctt	agggctgata	2340
gaggtacccc	cagatctcat	ggtccttatc	tctgactcag	cttaccaccag	aagaagaaga	2400
aaagcgaagg	gttcgcagag	agcggaaaca	gctggctgca	gctaagtgca	ggaaccgtcg	2460
gagggagctg	acagatcgac	ttcaggcggt	aaggaggagt	ctgggggtgt	cttgaggccg	2520
tgtggggagc	actctgcctt	gttcttcccc	cgtttctcac	tgtgcctgtg	tcctaaacga	2580
ggaaaccccc	tcttagggaa	caggggtcag	tataggetga	tggagtggct	ccatatgcat	2640
gctcagaccc	atgcccactt	actttcgact	gttccccact	ttccctgaat	atgtccccac	2700
atgtcacccct	cctggctttc	tctcagccta	aggagacaag	ctagaggagg	taattctctc	2760
accttctttt	cttcactaaa	taataatcca	ttttgccttc	ctgcctccat	ttttttttcc	2820
tgagctgggg	atctacctgt	cgtagttcag	ccctcctccc	ccaacttgat	agcctcaagt	2880
ttcagccctt	ggctgagatg	ccatcatcct	gactggctct	ggctggaaac	tattttgtgc	2940
taagtcaatt	ccttgtctgc	tacttcagct	atctacagty	ctgccgaact	tgagctgggtg	3000
gcgcccacca	agcccacttc	tttctctctt	ttttacctca	gtgcaacccc	ccacacacaa	3060
aacttcatgc	ctgccccttg	aaaccagggg	gcgtctctga	ctccccgtcg	ggaggctgaa	3120
ggagatgggt	aacagaacct	cattaaaaac	aacacataag	cattacctac	tgactcaaca	3180
aactgtagtg	tttttctttt	ttcctctcaa	aaaattattt	cgtttgttta	tttattattt	3240
gcttatgttt	gagtgagtgc	tgggtgcacca	cagcacacat	acgaggtcag	agggaaattt	3300
tcatagtttg	ttctctcctt	ccgtgttgtg	gggtgcttgc	ggcaatctcc	ttcactcagt	3360
gagctacaat	gcccccttct	gccctttaag	gcagagtact	ccttagtaca	gggggaccct	3420
ttcctcggcc	tctcaaagtt	gagattacaa	atgttcacca	tcacaccagg	cttggagttc	3480
ttgcctatca	gtgacgtcca	ctcctgccta	gcttcttccc	aaccatcttt	tagtctgatg	3540
gggaaaccga	ggcacgagta	gcattggtcta	ccaggatttc	ctcttagggg	acgggtcccct	3600
cagttgggag	ggagctgtcc	agccccctgg	atcagcagca	agaatgtatg	agtgtggggg	3660
tggggcgggtg	aagctactct	gtgtgggtcgc	tgaccagcaa	ttctcctttc	tctgtctcct	3720
atgacctggc	cctgctggga	tccattagga	aactgatcag	cttgaagagg	aaaaggcaga	3780
gctggagtgc	gagatcgccg	agctgcaaaa	agagaaggaa	cgcctggagt	ttgtcctggg	3840
ggcccacaaa	ccgggctgca	agatccccta	cgaagagggg	ccggggccag	gcccgtgggc	3900
cgaggtgaga	gatttgccag	ggtcaacatc	cgctaaggaa	gacggcttcg	gctggctgct	3960
gccgccccct	ccaccaccgc	ccctgccctt	ccagagcagc	cgagacgcac	cccccaacct	4020
gacggcttct	ctctttacac	acagtgaagt	tcaagtcttc	ggcgacccct	tccccgttgt	4080
tagcccttcg	tacacttctt	cgtttgtcct	cacctgcccc	gaggtctccg	cgttcgccgg	4140
cgcccaacgc	accagcggca	gcgagcagcc	gtccgacccg	ctgaactcgc	cctcccttct	4200
tgtctgtaa	actctttaga	caaacaaaac	aaacaaaccc	gcaaggaaca	aggaggagga	4260
agatgaggag	gagaggggag	gaagcagtc	gggggtgtgt	gtgtggacct	tttgactctt	4320
ctgtctgacc	acctgccgcc	tctgccatcg	gacatgacgg	aaggacctcc	tttgtgtttt	4380
gtgctctgtc	tctgggtttc	tgtgccccgg	cgagaccgga	gagctgggtga	ctttggggac	4440
aggggggtggg	gcggggatga	acacccctcc	tgcatactct	tgtcctgtta	cttcaaccca	4500
acttctgggg	atagatggct	gactgggtgg	gtaggggtggg	gtgcaacgcc	cacctttggc	4560
gtcttacgtg	aggctggagg	ggaaagagtg	ctgagtgtgg	gggtgcagggt	gggttgaggt	4620
cgagctggca	tgcacctcca	gagagaccca	acgaggaaat	gacagcaccg	tcctgtcctt	4680
cttttcccc	accaccccat	ccaccctcaa	gggtgcaggg	tgaccaagat	agctctgttt	4740
tgtcctctcg	ggccttagct	gattaactta	acatttccaa	gaggttacaa	cctcctcctg	4800
gacgaattga	gccccgact	gagggaaagtc	gatgccccct	ttgggagtct	gctaacccca	4860
cttcccgtg	attccaaaat	gtgaacccct	atctgactgc	tcagtctttc	cctcctggga	4920
aaactggctc	aggttggatt	tttttctctg	tctgctacag	agccccctcc	caactcaggc	4980
ccgtcccac	ccctgtgcag	tattatgcta	tgtccctctc	accctcaccc	ccacccagcg	5040
cgcccttggc	cgtcctcgtt	gggccttact	ggttttgggc	agcagggggc	gctgcgacgc	5100
ccatcttgct	ggagcgtttt	atactgtgaa	tgagtggctg	gattgctggg	cgcgccggat	5160
gggattgacc	cccagccctc	caaaactttt	cctgggcctc	cccttcttcc	acttgcttcc	5220
tccctcccc	tgacagggag	ttagactcga	aaggatgacc	acgacgcac	ccggtggcct	5280
tcttgctcag	gccccagact	ttttctcttt	aagtccttcg	ccttccccag	cctaggacgc	5340
caacttctcc	ccaccctggg	agccccgc	cctctcacag	aggctcaggc	aattttcaga	5400
gaagttttca	gggctgaggg	tttggctccc	ctatcctcga	tatttgaatc	cccaaatagt	5460
ttttggacta	gcatacttaa	gaggggggctg	agttcccact	atcccactcc	atccaattcc	5520
ttcagtccca	aagacgagtt	ctgtcccttc	cctccagctt	tcacctcgtg	agaatcccac	5580
gagtcagatt	tctattttct	aatattgggg	agatgggccc	taccgcccgt	ccccgtgct	5640
gcattggaaca	ttccataccc	tgtcctgggc	cctaggttcc	aaacctaate	ccaaacccca	5700



ccccagcta	tttatccctt	tcctgggtcc	caaaaagcac	ttatatctat	tatgtataaa	5760
taaatatatt	atatatgagt	gtgcgtgtgt	gtgcgtgtgc	gtgcgtgcgt	gcgtgcgtgc	5820
gagcttcctt	gttttcaagt	gtgctgtgga	gttcaaaatc	gcttctgggg	atgtgagtca	5880
gactttctgg	ctgtcccttt	ttgtcacttt	tttgttgttg	tctcggctcc	tctggctgtt	5940
ggagacagtc	ccggcctctc	cctttatcct	ttctcaagtc	tgtctcgctc	agaccacttc	6000
caacatgtct	ccactctcaa	tgactctgat	ctccggctctg	tctgttaatt	ctggatttgt	6060
cggggacatg	caattttact	tctgtaagta	agtgtgactg	ggtggtagat	tttttacaat	6120
ctatatcggt	gagaattctg	ggtggaaatg	tctgatcagg	agaagggcct	gccactgccg	6180
accacaatcc	attgactcca	tagccctcac	ccaggctgta	tttgtagatt	ttttcatttt	6240
gtttttttgt	attttgcacc	tgaccccggt	ggtgctgggg	cagtcctatca	ctgggcagct	6300
cccctccccc	ccttggttct	gcactgtcgc	caataaaaaa	cttttaaaaa	actgtatcct	6360
tcaggtcaaa	gtgtctgttt	tccttgga	tctactacat	ggcttccttt	cagaaaaacg	6420
gagtttggat	tgctagggaa	gtcttgctgg	cacttagtgg	gacgcctaac	gaatcagaac	6480
ctacaacggg	actaaaagga	agtggagact	tgctagggtt	tcccatgttc	ccaggctggg	6540
ccacctactt	gaaaaataa	ggggcgga	agtgtgaagt	accaaatttg	gtgaagggtc	6600
tgggagaatt	tcatgatcgg	aaaagaattt	attcaccttg	ggtgtgcaat	gaactttcag	6660
caacagttaa	gggcaagggt	gtaaaagctg	ggcacaactt	gtaaatccta	gcatttgaga	6720
ggtggaggca	aggggatcaa	ctggtggagt	tcagtgtcat	gtggatcgta	gataccaagc	6780
gcaaagatct	gctatgggga	gagggtctgg	tacaccaggg	gagccagaag	tttcgtggtg	6840
agggtagtgg	agggcaagtg	gagagtgaga	gttagcctca	gggagattct	acaggcaatg	6900
atgcagagtt	cagacgctcc	ctttgaaagc	actagagagc	cgcagcaggt	tttgagcaga	6960
gaaggttaga	gttaggtggg	ctcttctagc	ccatcccagg	ctgaggagga	cgctgagggt	7020
ttcaagaagg	atcgagaatg	gaaagcagag	gagaagaagg	atccaagagg	catggaggag	7080
gcagaacaca	tttctcttct	ttaatagcaa	gcctggaaag	gataacttgc	tgaggaggga	7140
gatgctcacc	agtcgggtgg	tctagggggg	tcttgga	gagaaggcat	ttgctcaagc	7200
ctcggttccc	ccattctcgc	tcttctgtca	gcttgtcttc	cattaagtgt	gtgtctcaag	7260
gccaccctgc	tcaggactcc	ttgtgagacg	accttctatg	ctcgagttca	ttaaaaacac	7320
aattgcctgg	tgccgtgctc	tctccactgg	ctcagttacc	tcaaaagacc	agggctaaag	7380
gtgtgatcac	aactctatcc	ccattactgc	tccaacgcag	agacaggact	gagccggagt	7440
gaacaaatga	acaaaaatga	ctaataatgc	atgcgtgatt	aaatacataa	aagagcagat	7500
gactggatga	gcaaatcggt	taaggagaga	cagcaagatc	ctagaatttt	ggagactaat	7560
ttaaatccat	ctttgagatg	catttggtcg	gaaattcctg	ggaggaaaaa	aagtgtaaat	7620
atgaagagag	aataaatgag	aataggggtg	gcttcagaga	ggttaactgc	gcgctggctg	7680
cttttgtaca	agaatgtgaa	ttgcaggggg	caaaatggga	tagatactcc	cgcccgaag	7740
gtggaattga	accactctgt	cgctaaacag	ctacagggtt	gaagcctgca	ccccagacca	7800
ctgaggatca	tccgggcgaa	aggagctatt	ttcagttagt	tatataaagg	cgagatacta	7860
ctacttttta	cacttatggg	cattatattgt	ggtatacagt	agataattaa	tttcaatggg	7920
ttcgaacatt	ttttttcact	ttttcttgtg	aacatgtgtt	tcctcagtaa	agtgttccgt	7980
gaatgactct	actaactaaa	aagtaagtag	cttcatttgc	atagcgctt	gcattttggg	8040
aagcagcgcc	taaagtgcct	gtctccctaa	ctaaaagcag	aattttttgc	aaagtga	8100
gtcagtttta	tttttgtttg	tttgtttgtc	tgtttgtttt	taatggaaaa	acttctcacg	8160
cggcccatte	gtagcagaat	tgcagatttt	ctgcaagcga	gaagcaagac	tttcgtaggg	8220
tctgacggca	cgcggccgca	gagcgacacc	tgccgttgct	ttatagaact	gcaagtatgt	8280
agggaatcta	ctgagtcctt	aggtgatgga	gttgacaacc	aactccctt	gagtttagac	8340
gctaaaaacc	atcccttttt	atatttatgt	gattagccca	gggaaactaa	ggctcagaca	8400
tggaataatac	cacagccgag	ttcttgtagc	ccaactccct	aggggaaatg	aaacctacag	8460
ttgtgggttt	aatatgcttg	gcccaggggc	agtggcccta	ttggcaggag	tgcccttatt	8520
agcggagggtg	taccttggtt	gagaagtgtg	tcacttggag	gcgaggtttt	gaggtacgta	8580
tgctcaagtc	tgccagtggt	gatcctgggt	gtctgcagaa	cgtgggtctc	ttctggctgc	8640
cttcggatca	aggtgtagaa	ctctcagctc	cttctccagc	accatgtctg	cctgcttaat	8700
gctttgcttc	tttccatgac	gataatgaac	tgtgctctg	aaactgtaag	tcagccccc	8760
agttacatgt	tttcttttat	aagagttgca	tatatatatg	tatgtatata	tgtatgtata	8820
tatgtatgta	tatatatata	tatatataaa	cagggtctca	ctcttttagct	ctggctggcc	8880
tgaaattcac	tatgtagccc	aggattgcct	gaacttgaa	gcaatcttcc	tgctcagcc	8940
tcccaatggg	attacaggca	tgagtcacaa	caagccattt	aaatcttatg	atgacttata	9000
agaagacaga	aatcagagt	tcctttacct	agttcacaga	tccttacaat	ctaacctcgt	9060
tcgctccata	aacagcccta	ccccaccctc	ctggaactgc	tttgagggaat	gctgcaggct	9120
ctcacaggca	cactcctcct	tggttaattct	cttcagcctg	gttgcccttc	ccccccatgt	9180
ccatgtggcc	caaagcctct	catcctgttc	tcaaatacca	ctagctagta	aggctccccg	9240
acctgaccgg	gtttaaatat	tagaaaaggg	tcactttctc	cctgccacag	acaaccaaac	9300
caccatatgc	ttgtcactta	ctacctgact	atgaagggtta	atagatgtct	tcacaacctt	9360
tctctgagcc	tcagtttccc	cacctgcata	atgcatctga	gacacagaat	tccttagagc	9420

tgtggttctc	ctcattccta	gtgctgggac	cctttaatac	atttcctcat	gttgtggtga	9480
ccccaccacc	accataaaat	tatttccatt	gatacttcat	aactgtaatt	ttttctattg	9540
ttatgaatag	taatgtaagc	atltgtgttt	cccagtgatc	ttagatgacc	ctgtggaaga	9600
gtcattccac	cccaaagggg	tccccaccac	aagttaagaa	ttcctgccat	agaggaatca	9660
cagggaccat	ggattaacac	ttgggtcgac	ttttgggctg	ccttctggga	ggcgctagag	9720
ctaatacag	ctacatcaat	ttctgaaatt	ttgtgtgtgt	gtgtgtgtgt	gtgtgtgtgt	9780
gtgtgtgtgt	gccctgagtc	gggtgctgag	ataggccagt	ggctttagt	ttcctggacc	9840
cattactcac	cagaactctc	ccctcacctg	attctttgat	gtgaacacta	tgtcttcata	9900
gtggcggtgg	caatagcagc	aacagtgaac	taaattttaa	aagtagaact	cagctggaga	9960
tacaaatatt	gcagttttga	agttgggggtg	gattgtctaa	taacttaata	acataaccca	10020
gaagagagggc	cccttggtct	tgcaaacttt	atatgcctca	gtacagggga	acgccagggc	10080
caagaagtgg	gagtgggtgg	gtaggggagc	agggtggggg	gagggtatag	gggactttcc	10140
ggatagcatt	tgaaatgtaa	atgaagaaaa	tatctaataa	aaatttgaaa	aaaaatgtta	10200
ccccagtttg	gcctggatct	cactacctca	accagactgg	catgtgactc	tgctgagatc	10260
tgctacttc	tgctcctgg	gtgcagaaga	caatttttgg	aagttagttc	tcttcttcca	10320
tcttgtggat	tccagggatt	gaactcgggt	catcaggctt	ggctgcaagt	gacttactta	10380
ggtgtctccc	agaccctctc	ggtttgatta	gttagatgct	gcacttcatg	cctgactttc	10440
gcactatgta	gatagagcaa	tgtctataac	atctcctaca	atgatatgta	tatcaagagc	10500
caagtgatga	gatggctcag	tgggtaagag	cacagactgc	tcttccaaag	gtcccagagt	10560
caaattcccag	caatcacata	gtggcttcca	ttccctctta	tggaatgtct	gaagactgct	10620
acagtgtact	tacatataat	aaataaataa	atcttaaaaa	aaaaaaacc	agccgggct	10680
ggtggcgcac	gcctttaatc	ccagcacttg	ggaggcagag	gcaggcggt	tcctgagttc	10740
gacgccagcc	tggtctacag	agtgagttcc	acgacagcca	gaactacaca	gagaaaccct	10800
gtctcgaaaa	aaaaaagaga	gagaggggag	tgagagcgca	ataatcttaa	catttctgtg	10860
gttgtctttg	ctgtagtcta	ttctgataag	caatgctggc	ttgctcccaa	ggtaggaagt	10920
aacatttctt	tataaaaggt	atltgctctg	ctttattttt	ctgttttatt	tatggtgctg	10980
aggatggaac	ccaggaccct	tggcaagcaa	ggctagctgt	ttaccactga	gccatactcc	11040
agccttgac	tgggggattc	taggcaaggg	ttctaccact	gagccacact	ccccacccc	11100
atccctctct	ggaagattct	aggcagttcc	atacctagcc	tttgatcttt	taagacgggt	11160
ttactagagc	tcagtt					11176

## INTERNATIONAL SEARCH REPORT

Int Application No  
PCT/US 99/30078A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N15/90 A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 30715 A (ISACOFF EHUD Y ;SIEGAL MICAH S (US); UNIV CALIFORNIA (US); CALIFOR) 16 July 1998 (1998-07-16) page 29, line 5 -page 30, line 10; claims ---	1-8,10, 22-26, 32,34,35
X	WO 98 28971 A (LINK CHRISTOPHER ;UNIV TECHNOLOGY CORP (US)) 9 July 1998 (1998-07-09) page 22, line 3 - line 6; figure 1  page 26, line 24 - line 28; claims 11-19; figure 1; example 3 page 18, line 9 - line 13 ---	1,2,8, 10,12, 14-17 1,2,7, 10-12, 14,17-20
Y	---	---

-/--

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*S\* document member of the same patent family

Date of the actual completion of the international search

26 June 2000

Date of mailing of the international search report

03/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3018

Authorized officer

Chambonnet, F



# INTERNATIONAL SEARCH REPORT

Int'l Application No  
PCT/US 99/30078

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>CONTAG CH, ET AL.: "Visualizing gene expression in living mammals using a bioluminescent"</p> <p>PHOTOCHEMISTRY AND PHOTOBIOLOGY, (1997)</p> <p>VOL. 66, NO. 4, PP. 523-531. ,</p> <p>XP000920570</p> <p>the whole document</p>	<p>1,2,7,</p> <p>10-12,</p> <p>14,17-20</p>
A	<p>CONTAG PR, OLOMU IN, STEVENSON DK, CONTAG CH. : "Bioluminescent indicators in living mammals."</p> <p>NAT MED. ,</p> <p>vol. 4, no. 2, February 1998 (1998-02),</p> <p>pages 245-247, XP000914588</p> <p>the whole document</p>	<p>1</p>
A	<p>WO 98 36081 A (ANGELIS DINO A DE ;SLOAN KETTERING INST CANCER (US); MIESENBOCK GE)</p> <p>20 August 1998 (1998-08-20)</p> <p>the whole document</p>	<p>1</p>
A	<p>WO 98 23633 A (CORNELL RES FOUNDATION INC)</p> <p>4 June 1998 (1998-06-04)</p> <p>the whole document</p>	<p>1</p>

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Int: I Application No

PCT/US 99/30078

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9830715 A	16-07-1998	AU 5090498 A	03-08-1998
WO 9828971 A	09-07-1998	AU 5619298 A	31-07-1998
		EP 0948612 A	13-10-1999
WO 9836081 A	20-08-1998	EP 0981633 A	01-03-2000
WO 9823633 A	04-06-1998	AU 7410398 A	22-06-1998

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**